

ORIGINAL RESEARCH ARTICLE

Faber polynomials estimates for bi-univalent functions of complex order involving q -derivative

Z. M. Saleh^{1,*}, A. O. Mostafa², S. M. Madian³

¹ Department of Basic Sciences, Higher Technological Institute, 10th of Ramadan 4955235, Egypt

² Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 2247358, Egypt

³ Department of Basic Sciences, Higher Institute of Engineering and Technology, New Damietta 7952567, Egypt

* Corresponding author: Z. M. Saleh, zeinabnsar2@gmail.com

ABSTRACT

In this paper, we define a new subclass of bi-univalent functions of complex order $\sum_q(\tau, \zeta; \varphi)$ which is defined by subordination in the open unit disc D by using $\nabla_q F(\vartheta)$ operator. Furthermore, using the Faber polynomial expansions, we get upper bounds for the coefficients of function belonging to this class. It is known that the calculus without the notion of limits is called q -calculus which has influenced many scientific fields due to its important applications. The generalization of derivative in q -calculus that is q -derivative was defined and studied by Jackson. A function $F \in A$ is said to be bi-univalent in D if both F and F^{-1} are univalent in D . The class consisting of bi-univalent functions is denoted by σ . The Faber polynomials play an important role in various areas of mathematical sciences, especially in geometric function theory. The purpose of our study is to obtain bounds for the general coefficients $|a_n| (n \geq 3)$ by using Faber polynomial expansion under certain conditions for analytic bi-univalent functions in subclass $\sum_q(\tau, \zeta; \varphi)$ and also, we obtain improvements on the bounds for the first two coefficients $|a_2|$ and $|a_3|$ of functions in this subclass. In certain cases, our estimates improve some of those existing coefficient bounds.

Keywords: analytic functions; bi-univalent functions; coefficient bounds; subordination; q -derivative; Faber polynomials
2010 Mathematical Subject Classification: 30C45

1. Introduction

Let A be the class of functions

$$F(\vartheta) = \vartheta + \sum_{\varepsilon=2}^{\infty} a_{\varepsilon} \vartheta^{\varepsilon}, \quad (1)$$

defined in $D = \{\vartheta \in \mathbb{C} : |\vartheta| < 1\}$ normalized by the conditions $F(0) = F'(0) - 1 = 0$ for every $\vartheta \in D$ and S be the subclass of A consisting of univalent functions in D . For every $F \in S$ there exists an inverse function F^{-1} which is defined in some neighborhood of the origin, and satisfying the conditions,

$$F^{-1}(F(\vartheta)) = \vartheta, (\vartheta \in D)$$

ARTICLE INFO

Received: 30 June 2023 | Accepted: 3 August 2023 | Available online: 22 August 2023

CITATION

Saleh ZM, Mostafa AO, Madian SM. Faber polynomials estimates for bi-univalent functions of complex order involving q -derivative. *Mathematics and Systems Science* 2023; 1(1): 2211. doi: 10.54517/mss.v1i1.2211

COPYRIGHT

Copyright © 2023 by author(s). *Mathematics and Systems Science* is published by Asia Pacific Academy of Science Pte. Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), permitting distribution and reproduction in any medium, provided the original work is cited.

and,

$$F^{-1}(F(\omega)) = \omega, (\vartheta < r_0(F); r_0(F) \geq \frac{1}{4}),$$

where,

$$g(\omega) = F^{-1}(\omega) = \omega - a_2\omega^2 + (2a_2^2 - a_3)\omega^3 + -(5a_2^3 - 5a_2a_3 + a_4)\omega^4 + \dots = \omega + \sum_{\varepsilon=2}^{\infty} A_{\varepsilon}\omega^{\varepsilon}. \quad (2)$$

If both F and F^{-1} are univalent in D , then $F \in \mathcal{A}$ is called bi-univalent in D and the class of these functions is denoted by σ . For there are many studies in this class^[1-6].

Faber^[7] introduced a polynomial which bears his name and is very important role in geometric function theory.

By using the expansion of this polynomial for $F \in \mathcal{S}$, the coefficients of its inverse $g = F^{-1}$ may be expressed as^[8,9]

$$g(\omega) = F^{-1}(\omega) = \omega + \sum_{\varepsilon=2}^{\infty} \frac{1}{\varepsilon} \chi_{\varepsilon-1}^{-\varepsilon}(a_2, a_3, \dots, a_{\varepsilon})\omega^{\varepsilon}, \quad (3)$$

where,

$$\begin{aligned} \chi_{\varepsilon-1}^{-\varepsilon} &= \frac{(-\varepsilon)!}{(-2\varepsilon+1)!(\varepsilon-1)!} a_2^{\varepsilon-1} + \frac{(-\varepsilon)!}{(2(-\varepsilon+1))!(\varepsilon-3)!} a_2^{\varepsilon-3} a_3 \\ &= \frac{(-\varepsilon)!}{(-2\varepsilon+3)!(\varepsilon-4)!} a_2^{\varepsilon-4} a_4 + \frac{(-\varepsilon)!}{(2(-\varepsilon+2))!(\varepsilon-5)!} a_2^{\varepsilon-5} \\ &\times [a_5 + (-\varepsilon+2)a_3^2] + \frac{(-\varepsilon)!}{(-2\varepsilon+5)!(\varepsilon-6)!} a_2^{\varepsilon-6} \times [a_6 + (-2\varepsilon+5)a_3a_4] \\ &+ \sum_{j \geq 7}^{\infty} a_2^{\varepsilon-j} V_j, \end{aligned}$$

such that V_j with $7 \leq j \leq \varepsilon$ is a homogeneous polynomial in the variables $a_2, a_3, \dots, a_{\varepsilon}$ ^[9]. The first three terms of $\chi_{\varepsilon-1}^{-\varepsilon}$ are

$$\chi_1^{-2} = -2a_2, \chi_2^{-3} = 3(2a_2^2 - a_3), \chi_3^{-4} = -4(5a_2^3 - 5a_2a_3 + a_4).$$

In general, for any $p \in \mathfrak{P} = \{0, \pm 1, \pm 2, \dots\}$, an expansion of χ_{ε}^p is

$$\chi_{\varepsilon}^p = p a_{\varepsilon+1} + \frac{p(p-1)}{2} H_{\varepsilon}^2 + \frac{p!}{(p-3)!3!} H_{\varepsilon}^3 + \dots + \frac{p!}{(p-\varepsilon)! \varepsilon!} H_{\varepsilon}^{\varepsilon},$$

where $H_{\varepsilon}^p = H_{\varepsilon}^p(a_2, a_3, \dots)$ and by Jahangiri et al.^[10], (see for details^[8,9,11,12,13])

$$H_{\varepsilon}^m(a_2, a_3, \dots, a_{\varepsilon+1}) = \sum_{\varepsilon=0}^{\infty} \frac{m! (a_2)^{\mu_1} \dots (a_{\varepsilon+1})^{\mu_{\varepsilon}}}{\mu_1! \dots \mu_{\varepsilon}!}, \quad (4)$$

where the sum is taken $\forall \mu_1, \dots, \mu_{\varepsilon} \in \mathbb{N} = \{1, 2, \dots\}$ satisfying

$$\begin{cases} \mu_1 + \mu_2 + \dots + \mu_{\varepsilon} = m, \\ \mu_1 + 2\mu_2 + \dots + k\mu_{\varepsilon} = \varepsilon. \end{cases}$$

Note that $H_{\varepsilon}^{\varepsilon}(a_2, a_3, \dots, a_{\varepsilon+1}) = a_2^{\varepsilon}$.

In the rest of this paper, assume that φ is an analytic function with positive real part in D , satisfying $\varphi(0) = 1$, $\varphi'(0) > 0$ and $\varphi(D)$ is symmetric w. r. to the real axis and has the expansion,

$$\varphi(\vartheta) = 1 + \psi_1\vartheta + \psi_2\vartheta^2 + \psi_3\vartheta^3 + \dots \quad (\psi_1 > 0).$$

Let $u(\vartheta)$ and $v(\vartheta)$ be analytic in D with $u(0) = v(0) = 0$, $|u(\vartheta)| < 1$, $|v(\vartheta)| < 1$, and

$$u(\vartheta) = \vartheta(p_1 + \sum_{\varepsilon=2}^{\infty} p_{\varepsilon}\vartheta^{\varepsilon-1}) \text{ and } v(\vartheta) = \vartheta(q_1 + \sum_{\varepsilon=2}^{\infty} q_{\varepsilon}\vartheta^{\varepsilon-1}) \quad (\vartheta \in D). \quad (5)$$

Then^[14],

$$p_1 \leq 1, \quad p_{\varepsilon} \leq 1 - p_1, \quad q_1 \leq 1, \quad q_{\varepsilon} \leq 1 - q_1, \quad (\varepsilon \in N \setminus \{1\}). \quad (6)$$

Jackson^[15] q – derivative, $0 < q < 1$, was defined by Annby and Mansour and other researchers^[16–21]:

$$\nabla_q F(\vartheta) = \begin{cases} \frac{F(\vartheta) - F(q\vartheta)}{(1-q)\vartheta}, & \vartheta \neq 0 \\ F'(0), & \vartheta = 0 \end{cases}$$

that is,

$$\nabla_q F(\vartheta) = 1 + \sum_{\varepsilon=2}^{\infty} [\varepsilon]_q a_{\varepsilon} \vartheta^{\varepsilon-1}, \quad (7)$$

where,

$$[j]_q = \frac{1 - q^j}{1 - q}, \quad [0]_q = 0. \quad (8)$$

As $q \rightarrow 1^-$, $[j]_q = j$ and $\nabla_q F(\vartheta) = F'(\vartheta)$.

Definition 1: For F, g , analytic in D , F is subordinate to g in D written $F < g$, if $\exists \Omega(\vartheta)$, analytic in D , with $\Omega(0) = 0$ and $|\Omega(\vartheta)| < 1$ ($\vartheta \in D$) such that $F(\vartheta) = g(\Omega(\vartheta))$ ($\vartheta \in D$)^[22,23].

Definition 2: For $\tau \in C^* = C \setminus \{0\}$, $0 \leq \zeta \leq 1$, $0 < q < 1$ and $F \in \sigma$, $F \in \Sigma_q(\tau, \zeta; \varphi)$ if for all $\vartheta, \omega \in D$ ∴

$$1 + \frac{1}{\tau} [\nabla_q(F(\vartheta)) + \zeta \vartheta \nabla_q(\nabla_q F(\vartheta)) - 1] < \varphi(\vartheta), \quad (9)$$

and

$$1 + \frac{1}{\tau} [\nabla_q(g(\omega)) + \zeta \omega \nabla_q(\nabla_q g(\omega)) - 1] < \varphi(\omega), \quad (10)$$

where $g(\omega) = F^{-1}(\omega)$.

Note that:

- 1) $\Sigma_q(1, \zeta; \phi) = \Sigma_q(\zeta; \phi)$;
- 2) $\Sigma_q((1 - \alpha)e^{-i\theta} \cos \theta, \zeta; \phi) = \Sigma_q(\zeta, \alpha, \theta; \phi)$, ($0 \leq \alpha < 1$, $|\theta| < \frac{\pi}{2}$), where

$$= \begin{cases} F \in \sigma : \frac{e^{i\theta} [\nabla_q(F(\vartheta)) + \zeta \vartheta \nabla_q(\nabla_q F(\vartheta))] - (\alpha \cos \theta + i \sin \theta)}{(1 - \alpha) \cos \theta} < \varphi(\vartheta) \\ g \in \sigma : \frac{e^{i\theta} [\nabla_q(g(\omega)) + \zeta \omega \nabla_q(\nabla_q g(\omega))] - (\alpha \cos \theta + i \sin \theta)}{(1 - \alpha) \cos \theta} < \varphi(\omega) \end{cases};$$
- 3) $\lim_{q \rightarrow 1^-} \Sigma_q(\tau, \zeta; \phi) = \Sigma(\tau, \zeta; \phi)$ ^[24];
- 4) $\lim_{q \rightarrow 1^-} \Sigma_q(1, \zeta; \phi) = \Sigma(\zeta; \phi)$ ^[25];
- 5) $\lim_{q \rightarrow 1^-} \Sigma_q((1 - \alpha)e^{-i\theta} \cos \theta, \zeta; \frac{1+\vartheta}{1-\vartheta}) = \Sigma(\zeta, \alpha, \theta; \frac{1+\vartheta}{1-\vartheta})$, ($0 \leq \alpha < 1$ $|\theta| < \frac{\pi}{2}$),

$$= \begin{cases} F \in \sigma : \frac{e^{i\theta}[F'(\vartheta) + \zeta\vartheta F''(\vartheta)] - (\alpha \cos \theta + i \sin \theta)}{(1 - \alpha) \cos \theta} < \frac{1 + \vartheta}{1 - \vartheta} \\ g \in \sigma : \frac{e^{i\theta}[g'(\omega) + \zeta\omega g''(\omega)] - (\alpha \cos \theta + i \sin \theta)}{(1 - \alpha) \cos \theta} < \frac{1 + \omega}{1 - \omega} \end{cases}$$

2. Main results

We assume that $\tau \in \mathbb{C}^*$, $0 < q < 1$, $0 \leq \zeta \leq 1$ and $F(\vartheta) \in \sigma$.

In this section we obtain some inequalities for the function class $\Sigma_q(\tau, \zeta; \phi)$.

Theorem 1: Let $\mathcal{F} \in \Sigma_q(\tau, \zeta; \phi)$. If $a_\epsilon = 0$ for $2 \leq \epsilon \leq \epsilon - 1$ then,

$$|a_\epsilon| \leq \frac{\psi_1 |\tau|}{(1 + \zeta[\epsilon - 1]_q) [\epsilon]_q} \quad (\epsilon \geq 3) \tag{11}$$

Proof: For functions $\nabla_q \mathcal{F}(\vartheta)$ given by Equation (7) and $g = F^{-1}$ we have:

$$\begin{aligned} & 1 + \frac{1}{\tau} [\nabla_q(\mathcal{F}(\vartheta)) + \zeta\vartheta \nabla_q(\nabla_q \mathcal{F}(\vartheta)) - 1] \\ &= 1 + \frac{1}{\tau} \sum_{\epsilon=2}^{\infty} (1 + \zeta[\epsilon - 1]_q) [\epsilon]_q a_\epsilon \vartheta^{\epsilon-1}, \end{aligned} \tag{12}$$

$$\begin{aligned} & 1 + \frac{1}{\tau} [\nabla_q(g(\omega)) + \zeta\omega \nabla_q(\nabla_q g(\omega)) - 1] \\ &= 1 + \frac{1}{\tau} \sum_{\epsilon=2}^{\infty} (1 + \zeta[\epsilon - 1]_q) [\epsilon]_q A_\epsilon \omega^{\epsilon-1}. \end{aligned} \tag{13}$$

Using Equation (3), we have:

$$\begin{aligned} & 1 + \frac{1}{\tau} [\nabla_q(g(\omega)) + \zeta\omega \nabla_q(\nabla_q g(\omega)) - 1] \\ &= 1 + \frac{1}{\tau} \sum_{\epsilon=2}^{\infty} (1 + \zeta[\epsilon - 1]_q) [\epsilon]_q \frac{1}{\epsilon} \chi_{\epsilon-1}^{-\epsilon}(a_2, a_3, \dots, a_\epsilon) \omega^{\epsilon-1}. \end{aligned} \tag{14}$$

Considering Equations (9) and (10), there are two Schwarz functions $u, v : \mathbb{D} \rightarrow \mathbb{D}$ with $u(0) = v(0) = 0$, which are given by Equation (5), so that,

$$1 + \frac{1}{\tau} [\nabla_q(F(\vartheta)) + \zeta\vartheta \nabla_q(\nabla_q F(\vartheta)) - 1] = \phi(u(\vartheta)), \tag{15}$$

$$1 + \frac{1}{\tau} [\nabla_q(g(\omega)) + \zeta\omega \nabla_q(\nabla_q g(\omega)) - 1] = \phi(v(\omega)). \tag{16}$$

Also, by Equation (4) we get:

$$\begin{aligned} \phi(u(\vartheta)) &= 1 + \psi_1 p_1 \vartheta + (\psi_1 p_2 + \psi_2 p_1^2) \vartheta^2 + \dots \\ &= 1 + \sum_{\epsilon=1}^{\infty} \sum_{\epsilon=1}^{\epsilon} \psi_\epsilon D_\epsilon^\epsilon(p_1, p_2, \dots, p_\epsilon) \vartheta^\epsilon \quad (\vartheta \in \mathbb{D}), \end{aligned} \tag{17}$$

and,

$$\begin{aligned} \phi(v(\omega)) &= 1 + \psi_1 q_1 \omega + (\psi_1 q_2 + \psi_2 q_1^2) \omega^2 + \dots \\ &= 1 + \sum_{\epsilon=1}^{\infty} \sum_{\epsilon=1}^{\epsilon} \psi_{\epsilon} D_{\epsilon}^{\epsilon}(q_1, q_2, \dots, q_{\epsilon}) \omega^{\epsilon} \quad (\omega \in \mathbb{D}). \end{aligned} \tag{18}$$

Comparing the coefficients of Equations (12), (15) and (17), we get:

$$\frac{1}{\tau} (1 + \zeta[\epsilon - 1]_q) [\epsilon]_q a_{\epsilon} = \sum_{l=1}^{\epsilon-1} \psi_l H_{\epsilon-1}^l(p_1, p_2, \dots, p_{\epsilon-1}) \quad (\epsilon \geq 2). \tag{19}$$

Similarly, from Equations (14), (16) and (18), we get:

$$\frac{1}{\tau} (1 + \zeta[\epsilon - 1]_q) [\epsilon]_q \frac{1}{\epsilon} \chi_{\epsilon-1}^{-\epsilon}(a_2, a_3, \dots, a_{\epsilon}) = \sum_{\epsilon=1}^{\epsilon-1} \psi_{\epsilon} D_{\epsilon-1}^{\epsilon}(q_1, q_2, \dots, q_{\epsilon-1}) \quad (\epsilon \geq 2). \tag{20}$$

Now, from $a_{\epsilon} = 0$ for $2 \leq \epsilon \leq \epsilon - 1$, we have $A_{\epsilon} = -a_{\epsilon}$ and the Equations (19) and (20) yield

$$\begin{aligned} (1 + \zeta[\epsilon - 1]_q) [\epsilon]_q a_{\epsilon} &= \tau \psi_1 p_{\epsilon-1}, \\ - (1 + \zeta[\epsilon - 1]_q) [\epsilon]_q a_{\epsilon} &= \tau \psi_1 q_{\epsilon-1}. \end{aligned} \tag{21}$$

Taking the modulus of each of the two equations in Equation (21) and using Equation (6), we obtain Equation (11).

Corollary 1: For $\varphi(\vartheta) = \left(\frac{1+\vartheta}{1-\vartheta}\right)^{\alpha}$ ($0 < \alpha \leq 1$), let $\mathcal{F} \in \Sigma_q(\tau, \zeta; \phi)$ then

$$|a_{\epsilon}| \leq \frac{2\alpha |\tau|}{(1 + \zeta[\epsilon - 1]_q) [\epsilon]_q} \quad (\epsilon \geq 3). \tag{22}$$

Corollary 2: For $\varphi(\vartheta) = \frac{1+(1-2\beta)\vartheta}{1-\vartheta}$ ($0 \leq \beta < 1$), let $\mathcal{F} \in \Sigma_q(\tau, \zeta; \phi)$ then

$$|a_{\epsilon}| \leq \frac{2 |\tau| (1 - \beta)}{(1 + \zeta[\epsilon - 1]_q) [\epsilon]_q} \quad (\epsilon \geq 3). \tag{23}$$

Remark 1: For $\tau = 1$, $q \rightarrow 1^-$ in Corollary 2, we get results obtained by Srivastava et al.^[13], for all $0 \leq \zeta \leq 1$.

Theorem 2: Let $\mathcal{F} \in \Sigma_q(\tau, \zeta; \phi)$. Then,

$$|a_2| \leq \frac{\psi_1 \sqrt{\psi_1} |\tau|}{\sqrt{\psi_1 [2]_q^2 (1 + \zeta)^2 + \left| \tau [3]_q (1 + \zeta [2]_q) \psi_1^2 - [2]_q^2 (1 + \zeta)^2 \psi_2 \right|}}, \tag{24}$$

$$|a_3| \leq \min \{ \mathcal{K}(\zeta), \mathcal{L}(\zeta) \}, \tag{25}$$

where,

$$\mathcal{L}(\zeta) = \begin{cases} \frac{\psi_1 |\tau|}{[3]_q (1 + \zeta [2]_q)} \times \frac{[3]_q (1 + \zeta [2]_q) |\tau| \psi_1^2 + [3]_q (1 + \zeta [2]_q) \tau \psi_1^2 - [2]_q^2 (1 + \zeta)^2 \psi_2}{[2]_q^2 (1 + \zeta)^2 \psi_1 + [3]_q (1 + \zeta [2]_q) \tau \psi_1^2 - [2]_q^2 (1 + \zeta)^2 \psi_2}, & \psi_1 \geq \frac{[2]_q^2 (1 + \zeta)^2}{[3]_q (1 + \zeta [2]_q) |\tau|} \\ \frac{\psi_1 |\tau|}{[3]_q (1 + \zeta [2]_q)}, & 0 \leq \psi_1 \leq \frac{[2]_q^2 (1 + \zeta)^2}{[3]_q (1 + \zeta [2]_q) |\tau|} \end{cases}, \tag{26}$$

and

$$\mathcal{K}(\zeta) = \begin{cases} \frac{|\psi_2|\tau}{[3]_q(1+\zeta[2]_q)} & , |\psi_2| > \psi_1 \\ \frac{\psi_1\tau}{[3]_q(1+\zeta[2]_q)} & , |\psi_2| \leq \psi_1 \end{cases} . \quad (27)$$

Proof: If we set $\varepsilon = 2$ and $\varepsilon = 3$ in Equations (19) and (20), respectively, we have

$$\frac{1}{\tau} [2]_q(1 + \zeta)a_2 = \psi_1 p_1, \quad (28)$$

$$\frac{1}{\tau} [3]_q(1 + \zeta[2]_q)a_3 = \psi_1 p_2 + \psi_2 p_1^2, \quad (29)$$

$$-\frac{1}{\tau} [2]_q(1 + \zeta)a_2 = \psi_1 q_1, \quad (30)$$

and

$$\frac{1}{\tau} [3]_q(1 + \zeta[2]_q)(2a_2^2 - a_3) = \psi_1 q_2 + \psi_2 q_1^2. \quad (31)$$

From Equations (28) and (30), we obtain:

$$p_1 = -q_1. \quad (32)$$

Adding Equations (29) and (31), and using Equation (32), we have:

$$\frac{2}{\tau} [3]_q(1 + \zeta[2]_q)a_2^2 - 2p_1^2\psi_2 = \psi_1(p_2 + q_2). \quad (33)$$

From Equation (28), we get:

$$\left[2\tau[3]_q\psi_1^2(1 + \zeta[2]_q) - 2[2]_q^2(1 + \zeta)^2\psi_2 \right] a_2^2 = \tau^2\psi_1^3(p_2 + q_2). \quad (34)$$

By Equations (6), (28) and (32), we obtain:

$$\begin{aligned} & \left| 2\tau[3]_q\psi_1^2(1 + \zeta[2]_q) - 2[2]_q^2(1 + \zeta)^2\psi_2 \right| |a_2|^2 \\ & \leq |\tau|^2 \psi_1^3(|p_2| + |q_2|) \\ & \leq 2|\tau|^2 \psi_1^3(1 - |p_1|^2) \\ & = 2|\tau|^2 \psi_1^3 - 2[2]_q^2(1 + \zeta)^2\psi_1 |a_2|^2 . \end{aligned} \quad (35)$$

Consequently,

$$|a_2|^2 \leq \frac{|\tau|^2 \psi_1^3}{[2]_q^2(1 + \zeta)^2\psi_1 + \left| \tau[3]_q\psi_1^2(1 + \zeta[2]_q) - [2]_q^2(1 + \zeta)^2\psi_2 \right|}.$$

So, we obtain the bound on $|a_2|$ in Equation (24).

Next, in order to find the bound on the coefficient $|a_3|$, by subtracting Equation (31) from Equation (29), and using Equation (32), we get:

$$\frac{2}{\tau} [3]_q(1 + \zeta[2]_q)a_2^2 - \frac{2}{\tau} [3]_q(1 + \zeta[2]_q)a_3 = \psi_1(q_2 - p_2). \quad (36)$$

Using Equation (6), we have:

$$\begin{aligned} 2[3]_q(1 + \zeta[2]_q)|a_3| & \leq 2[3]_q(1 + \zeta[2]_q)|a_2|^2 + 2|\tau|\psi_1(|p_2| + |q_2|) \\ & \leq 2[3]_q(1 + \zeta[2]_q)|a_2|^2 + 2|\tau|\psi_1(1 - |p_1|^2). \end{aligned} \quad (37)$$

From Equation (28), we get:

$$[3]_q (1 + \zeta [2]_q) |\tau| \psi_1 |a_3| \leq |\tau|^2 \psi_1^2 + \left[|\tau| [3]_q \psi_1 (1 + \zeta [2]_q) - [2]_q^2 (1 + \zeta)^2 \right] |a_2|^2. \quad (38)$$

On the other hand, from Equation (29), we have:

$$[3]_q (1 + \zeta [2]_q) |a_3| \leq \tau \left[\psi_1 (1 - |p_1|^2) + |\psi_2| |p_1|^2 \right].$$

Consequently,

$$|a_3| \leq \begin{cases} \frac{|\psi_2| \tau}{[3]_q (1 + \zeta [2]_q)} & , |\psi_2| > \psi_1 \\ \frac{\psi_1 \tau}{[3]_q (1 + \zeta [2]_q)} & , |\psi_2| \leq \psi_1 \end{cases}. \quad (39)$$

Hence, from Equations (38) and (39), we obtain Equation (25).

By letting $\zeta = 0$, $\tau = 1$, we have:

Corollary 3: Let $\mathcal{F} \in \sum_q(1, 0; \phi)$. Then

$$|a_3| \leq \min \{ \mathcal{K}(0), \mathcal{L}(0) \}, \quad (40)$$

where,

$$\mathcal{L}(0) = \begin{cases} \frac{\psi_1}{[3]_q} \times \frac{[3]_q \psi_1^2 + |[3]_q \psi_1^2 - [2]_q^2 \psi_2|}{[2]_q^2 \psi_1 + |[3]_q \psi_1^2 - [2]_q^2 \psi_2|} & , \psi_1 \geq \frac{[2]_q^2}{[3]_q} \\ \frac{\psi_1}{[3]_q} & , 0 \leq \psi_1 \leq \frac{[2]_q^2}{[3]_q} \end{cases}, \quad (41)$$

and,

$$\mathcal{K}(0) = \begin{cases} \frac{|\psi_2|}{[3]_q} & , |\psi_2| > \psi_1 \\ \frac{\psi_1}{[3]_q} & , |\psi_2| \leq \psi_1 \end{cases}. \quad (42)$$

3. Future work

The authors suggest to find upper bounds for the coefficients of function class $\sum_{\lambda,q}^m(\tau, \zeta; \phi)$ for all $\vartheta, \omega \in D$:

$$1 + \frac{1}{\tau} \left[\nabla_{\lambda,q}^m(F(\vartheta)) + \zeta \vartheta \nabla_q(\nabla_{\lambda,q}^m F(\vartheta)) - 1 \right] < \varphi(\vartheta), \quad (43)$$

and,

$$1 + \frac{1}{\tau} \left[\nabla_{\lambda,q}^m(g(\omega)) + \zeta \omega \nabla_q(\nabla_{\lambda,q}^m g(\omega)) - 1 \right] < \varphi(\omega), \quad (44)$$

where,

$$\nabla_{\lambda,q}^m(F(\vartheta)) = \vartheta + \sum_{k=2}^{\infty} [1 + \lambda([k]_q - 1)]^m a_k \vartheta^k, \lambda \geq 0, m \in N_0 = N \cup \{0\}, \quad (45)$$

is the q – Al-Oboudi operator is defined by Aouf et al.^[26].

4. Conclusions

Throughout the paper, we defined a new subclass of bi-univalent functions of complex order $\sum_q(\tau, \zeta; \phi)$ by using $\nabla_q F(\vartheta)$ operator. Furthermore, using the Faber polynomial expansions, we find the initial coefficient bounds for this function class. This paper generalized many results.

Author contributions

Conceptualization, AOM and SMM; methodology, AOM and SMM; validation, ZMS, AOM and SMM; formal analysis, ZMS, AOM and SMM; investigation, ZMS, AOM and SMM; resources, AOM and SMM; data curation, ZMS; writing—original draft preparation, ZMS; writing—review and editing, AOM and SMM; supervision, AOM and SMM. All authors have read and agreed to the published version of the manuscript.

Conflict of interest

The authors declare no conflict of interest.

References

1. Ali RM, Lee SK, Ravichandran V, Supramanian S. Coefficient estimates for bi-univalent Ma-Minda starlike and convex functions. *Applied Mathematics Letters* 2012; 25(3): 344–351. doi: 10.1016/j.aml.2011.09.012
2. Aouf MK, Madian SM. Coefficient bounds for bi-univalent classes defined by Bazilevič functions and convolution. *Boletín de la Sociedad Matemática Mexicana* 2020; 26(2): 1045–1062. doi: 10.1007/s40590-020-00304-0.
3. Bulut S, Magesh N, Balaji VK. Initial bounds for analytic and bi-univalent functions by means of Chebyshev polynomials. *Journal of Classical Analysis* 2017; 11(1): 83–89. doi: 10.7153/jca-2017-11-06
4. Madian SM. Some properties for certain class of bi-univalent functions defined by q -Cătaș operator with bounded boundary rotation. *AIMS Mathematics* 2022; 7(1): 903–914. doi: 10.3934/math.2022053
5. Mostafa AO, Aouf MK, Elmorsy RE. Coefficient bounds for general class of bi-univalent functions of complex order associated with q -Sălăgean operator and Chebyshev polynomials. *Electronic Journal of Mathematical Analysis and Applications* 2020; 8(2): 251–260.
6. Mostafa AO, Saleh ZM. Coefficient bounds for a class of bi-univalent functions defined by chebyshev polynomials. *International Journal of Open Problems in Complex Analysis* 2021; 13(3): 19–28.
7. Faber G. Über polynomische Entwicklungen II. *Mathematische Annalen* 1903; 57(3): 389–408. doi: 10.1007/BF01449884
8. Airault H, Bouali A. Differential calculus on the Faber polynomials. *Bulletin des Sciences Mathématiques* 2006; 130(3): 179–222. doi: 10.1016/j.bulsci.2005.10.002
9. Airault H, Ren J. An algebra of differential operators and generating functions on the set of univalent functions. *Bulletin des Sciences Mathématiques* 2002; 126(5): 343–367. doi: 10.1016/S0007-4497(02)01115-6
10. Jahangiri JM, Hamidi SG. Coefficient estimates for certain classes of bi-univalent functions. *International Journal of Mathematics and Mathematical Sciences* 2013; 2013: 190560. doi: 10.1155/2013/190560
11. Todorov PG. On the Faber polynomials of the univalent functions of class Σ . *Journal of Mathematical Analysis and Applications* 1991; 162(1): 268–276. doi: 10.1016/0022-247X(91)90193-4
12. Airault H. Symmetric sums associated to the factorization of Grunsky coefficients. In: *Groups and Symmetries: From Neolithic Scots to John McKay*. CRM Proceedings and Lecture Notes; 2009.
13. Srivastava HM, Eker SS, Ali RM. Coefficient bounds for a certain class of analytic and bi-univalent functions. *Filomat* 2015; 29(8): 1839–1845. doi: 10.2298/FIL1508839S
14. Nehari Z. *Conformal Mapping*. McGraw-Hill Book; 1952.
15. Jackson FH. On q -functions and a certain difference operator. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* 1909; 46(2): 253–281. doi: 10.1017/S0080456800002751
16. Annby MH, Mansour ZS. *q -Fractional Calculus Equations*. Springer; 2012.
17. Aouf MK, Mostafa AO. Subordination results for analytic functions associated with fractional q -calculus operators with complex order. *Afrika Matematika* 2020; 31(2): 1387–1396. doi: 10.1007/s13370-020-00803-3
18. Aouf MK, Mostafa AO. Some subordinating results for classes of functions defined by Sălăgean type q derivative operator. *Filomat* 2020; 34(7): 2283–2292. doi: 10.2298/FIL2007283A
19. Aral A, Gupta V, Agarwal RP. *Applications of q -Calculus in Operator Theory*. Springer; 2013.
20. Frasin BA, Murugusundaramoorthy G. A subordination results for a class of analytic functions defined by q -differential operator. *Annales Universitatis Paedagogicae Cracoviensis Studia Mathematica* 2020; 19(1): 53–64. doi: 10.2478/aupcsm-2020-0005
21. Srivastava HM, Aouf MK, Mostafa AO. Some properties of analytic functions associated with fractional q -calculus operators. *Miskolc Mathematical Notes* 2019; 20(2): 1245–1260. doi: 10.18514/MMN.2019.3046
22. Bulboacă T. Differential subordinations and superordinations-preserving integral operators. Available online: <https://www.researchgate.net/profile/Teodor-Bulboaca->

2/publication/266915052_Differential_subordination_and_superordination-preserving_integral_operators/links/565d88d308aeafc2aac7e3fd/Differential-subordination-and-superordination-preserving-integral-operators.pdf (accessed on 2 May 2023).

23. Miller SS, MocanuPT. *Differential Subordinations—Theory and Applications*. CRC Press; 2000.
24. Deniz E. Certain subclasses of bi-univalent functions satisfying subordinate conditions. *Journal of Classical Analysis* 2013; 2(1): 49–60. doi: 10.7153/jca-02-05
25. Adegani EA, Bulut S, Zireh A. Coefficient estimates for a subclass of analytic bi-univalent functions. *Bulletin of the Korean Mathematical Society* 2018; 55(2): 405–413. doi: 10.4134/BKMS.b170051
26. Aouf MK, Mostafa AO, Elmorsy RE. Certain subclasses of analytic functions with varying arguments associated with q -difference operator. *Afrika Matematika* 2021; 32: 621–630. doi: 10.1007/s13370-020-00849-3