

# Effects of fertilizer reduction on Yield and non-point source pollution of Yongyou rice

Maoqian, Zhoujiangming

(Jiangshan Agricultural Technology Extension Center, jiangshan 324100, zhejiang)

man\_0034@163.com

## Abstract:

In the main grain producing areas of China, there are still problems of agricultural non-point source pollution and eutrophication of surrounding surface water caused by excessive application of chemical fertilizers. In order to change the unreasonable traditional fertilization habits of farmers and minimize agricultural non-point source pollution, a fertilizer reduction fertilization project was carried out in Jiangshan City to study the effects of 10%~30% fertilizer reduction on the yield of rice Yongyou 15 and the loss of nutrients (total nitrogen, total phosphorus and total potassium) in surface runoff of paddy fields on the basis of farmers' conventional fertilization. The results showed that there was no significant change in rice yield by reducing the amount of conventional fertilizer by 10%~20%, and the yield by reducing 10% was the highest. The nutrient concentration in the drainage of paddy field quickly rose to the highest value 1h after fertilization, and then nitrogen, phosphorus and potassium quickly decreased by 25.9%~66.0%, 70.1%~88.3% and 25.0%~52.5% after 24h. After that, all nutrients decreased slowly until the end of the experiment. This shows that if the paddy field irrigation water outflow is caused by heavy rain or artificial drainage, the nutrient diversion loss risk period (also a high risk period of non-point source pollution) is within a few days after fertilization. The reduction of chemical fertilizer has a significant impact on the nutrient content in the drainage of paddy field, and the amount of conventional fertilization is reduced by 10%~30% one hour after fertilization, the contents of total nitrogen, total phosphorus and total potassium in the drainage decreased by 3.7%~68.2%, 26.3%~64.8% and 5.8%~57.5% respectively. It is of great significance to improve economic benefits and protect the ecological environment in rice production.

**Keywords:** Reducing fertilization; Yongyou 15; rice; Nutrient loss; Non point source pollution

Rice is one of the main food crops in China, with a rice area of  $2.969 \times 10^7 \text{hm}^2$ , total output  $2.096 \times 10^{11} \text{kg}$ <sup>[1]</sup>, with the area and output ranking second after that of corn, providing a basic guarantee for China's food security. As a basic element of agricultural production, chemical fertilizer has played an important role in increasing rice yield <sup>[2-3]</sup>; however, while chemical fertilizer has greatly

increased crop yield, it has also led to people's dependence on it, which not only leads to the phenomenon of increasing fertilizer without increasing production, but also pollutes the air, soil and water, bringing huge environmental pressure to the sustainable development of agriculture <sup>[3-4]</sup>. Makun et al. <sup>[2]</sup> analyzed the contribution of chemical fertilizer to grain production by using the panel data model. They

---

Fund Project: Fertilizer reduction and efficiency increase project of the Ministry of agriculture and rural areas

About the author: Maoqian (1992 -), female, Jiangshan, Zhejiang Province, assistant agronomist, bachelor degree, mainly engaged in soil and fertilizer, e-mail: 1592560282@qq. Com.

Corresponding author: Zhoujiangming, promotion researcher, e-mail: man\_0034@163. Com.

believed that since 2000, china's chemical fertilizer input has exceeded the optimal application amount in economics and entered the stage of diminishing marginal return. Under the condition that the input of other factors is relatively stable, increasing the amount of chemical fertilizer will not bring about the expected increase in production, but will increase the production cost and increase the risk of polluting the agricultural production environment. They suggested that the amount of chemical fertilizer applied should be appropriately reduced.

In recent years, many technicians have carried out research on the impact of fertilizer reduction on rice production and environmental pollution, and achieved great results. Lihongjiang et al. [5] conducted a reduction test on the customary nitrogen application amount. The results showed that under the condition of reducing 10% nitrogen fertilizer, the rice yield was not affected and the total nitrogen loss rate of surface runoff decreased by 9.2%; shenjianguo et al. [6] applied different slow-release fertilizers to reduce the yield by 20% and 25%, but the rice yield increased by 8.8% and 5.0%. However, there are relatively few studies on the effect of fertilizer reduction on nitrogen and phosphorus in paddy field drainage at the same time. Therefore, under the condition of conventional fertilization, this paper studied the effect of fertilizer reduction treatment on the nitrogen and phosphorus content in the field water layer and rice yield, in order to optimize the scientific fertilization technology of reducing non-point source pollution without rice yield reduction, and provide technical support for improving the agricultural ecological environment.

## 1 Materials and methods

### 1.1 Test materials

The experiment was conducted in the rice field of Jianai family farm, huqian village, hecun Town, jiangshan City. The test site has a typical subtropical monsoon climate, and the agricultural production is mainly rice planting. The soil of the test field is slightly acidic, with clay texture and high content of organic matter. The specific physical and chemical properties are total nitrogen  $14\text{g}\cdot\text{kg}^{-1}$ , available phosphorus  $224.1\text{mg}\cdot\text{kg}^{-1}$ , available potassium  $48.0\text{mg}\cdot\text{kg}^{-1}$ , organic matter  $27.1\text{g}\cdot\text{kg}^{-1}$  and  $\text{pH}5.42$ .

The tested rice variety was Yongyou 15. The fertilizer is 45% compound fertilizer (N15%,  $\text{P}_2\text{O}_5$ 15%,  $\text{K}_2\text{O}$ 15%), urea (N46%) and potassium chloride ( $\text{K}_2\text{O}$ 60%).

### 1.2 Treatment design

There were 5 treatments: no fertilization (CK); conventional fertilization (CF) for farmers:  $\text{N}210\text{kg}\cdot\text{hm}^{-2}$ ,  $\text{P}_2\text{O}_590\text{kg}\cdot\text{hm}^{-2}$ ,  $\text{K}_2\text{O}150\text{kg}\cdot\text{hm}^{-2}$ ; farmers' conventional fertilization and chemical fertilizer shall be reduced by 10% (90%CF); 20% reduction of conventional fertilization and chemical fertilizer for farmers (80%CF); farmers' conventional fertilization and chemical fertilizer shall be reduced by 30% (70%CF). The small area is  $35\text{m}^2$ , repeated for 3 times, and arranged in random blocks.  $30\text{cm}\times 30\text{cm}$  wide for inter district use. The high ridge is separated by independent irrigation and drainage, and the ridge is covered with black film to avoid mutual leakage of water and fertilizer between communities. Protective rows are set around.

Rice will be sown on June 5, 2020, transplanted on July 1, and matured and harvested on October 30. The base fertilizer was applied on the test field on June 28, and the amount of N,  $\text{P}_2\text{O}_5$  and  $\text{K}_2\text{O}$  fertilizer accounted for 26.7%, 62.5% and 37.5% of the total fertilizer respectively. After the base fertilizer was applied to each plot, the iron rake was used

to manually mix the fertilizer into the arable soil, and the rest of the fertilizer was directly applied to the surface of the paddy field as Topdressing on July 9. The management of irrigation water and pest control in each community shall be carried out in a unified manner with reference to the habits of local farmers.

### 1.3 Investigation and analysis

Before the test, a soil sample was collected from the paddy field to represent the test soil properties and analyze the soil total nitrogen, available phosphorus, available potassium, organic matter and pH. At the mature stage of rice, the yield of each plot is calculated by actual cutting. After the rice is dried in the sun, the water content is measured, and finally converted into the rice yield with 14% water content.

During the experiment, the flooded layer of about 2cm was maintained within one week after transplanting. After the application of base fertilizer and top dressing, the paddy field ponding samples were collected. After the drainage outlet was drained for 10min, 1kg of irrigation water was collected and put into polyethylene bottles for sealing in time. Three samples were collected repeatedly in each plot. At first, water samples were collected at 1 and 4 hours after fertilization, and then samples were collected every day for 5 consecutive days (not collected on the fourth day after topdressing). A total of 7 water samples were collected after base fertilizer and 6 water samples were collected after topdressing. All samples shall be sent to the local laboratory for analysis of total nitrogen, total phosphorus and total potassium. The samples were digested with  $K_2S_2O_8$ -NaOH and analyzed by UV spectrophotometry and flame photometer respectively. The nutrient loss of surface runoff is calculated by multiplying the volume of paddy water by the nutrient depth of the day.


### 1.4 Statistical methods

The average yield of rice among different treatments and the nutrient content of water samples at the same time were statistically analyzed by DPS9.5, multiple comparisons were made by LSD method, and the significance was tested at the level of  $p < 0.05$ . The correlation diagram was made by Origin9.0 software.

## 2 Results and analysis

### 2.1 Output

Figure 1 shows that the yield of fertilizer treatment is 4.0%~13.4% higher than that of no fertilizer treatment. Among them, 90%CF treatment had the highest yield, which was  $6863.5 \text{ kg} \cdot \text{hm}^{-2}$ , significantly higher than CK and 70%CF treatment, and had no significant difference with CF and 80%CF. The results showed that the conventional fertilization amount of farmers was high, while the reduction of chemical fertilizer by more than 30% was low, which had a negative impact on rice yield, while the reduction of 10%~20% had no significant impact on rice yield, especially the reduction of chemical fertilizer by 10% was the best amount of chemical fertilizer.



There was no same small letter between columns, indicating significant difference between groups ( $p < 0.05$ ).

Yield
Handle

Fig. 1 Rice yield of different treated water


## 2.2 Field water nutrients

As shown in Figure 2, the nutrient content in the field ponding increased rapidly after deep application of base fertilizer, and the total nitrogen and total potassium content reached the peak at 4h, which were 61.6 and 64.6mg·L<sup>-1</sup> respectively, 9.7 and 6.6 times that of no fertilizer (CK), with significant difference. At 24h, the content remained high (60.1 and 63.3mg·L<sup>-1</sup>). Two days after fertilization, there was a heavy rainstorm (Fig. 3). From June 30 to July 1, the average daily rainfall reached more than 150mm. All irrigation ponding in the test field was discharged through surface runoff, and the contents of total nitrogen and total potassium in the remaining ponding rapidly decreased to CK level. At 72h, the contents of total nitrogen and total potassium in fertilization treatment increased, and were significantly higher than CK, which were 3.6 times and 2.9 times of CK. Then it decreased gradually, and there was no significant difference with CK.

The change of total phosphorus content in paddy field ponding was different from that of nitrogen and potassium. Total phosphorus reached its peak at 2.4mg·L<sup>-1</sup> after 1h of fertilization, and rapidly decreased to 0.5mg·L<sup>-1</sup> after 4h. After 24h, it was basically the same as CK. The above results showed that the content of nitrogen and potassium in the paddy field ponding remained high after applying base fertilizer, and decreased and stabilized at a low level after at least 96h, while the content of phosphorus in the paddy field ponding rapidly decreased to the level of the unfertilized paddy field after reaching the peak value about 1h, and the content of phosphorus in the paddy field ponding was extremely unstable.

Figure 4 shows that the chemical fertilizer has been basically dissolved 1h after topdressing,

and the contents of total nitrogen, total phosphorus and total potassium in the accumulated water on the field surface have rapidly increased to the peak (nitrogen 269.7mg·L<sup>-1</sup>, phosphorus 72.2mg·L<sup>-1</sup>, potassium 98.0mg·L<sup>-1</sup>), and then gradually decreased. Among them, the total nitrogen content in ponding decreased significantly after 24 hours of fertilization. Compared with 1 hour after fertilization, the total nitrogen content in each fertilization treatment decreased by 25.9%~66.0%, and the decline increased with the increase of fertilizer dosage, and then decreased slowly. Compared with the total nitrogen content of CF, there was no significant difference between 90%CF and 80%CF within 96h after fertilization, while 70%CF was significantly lower than that of CF treatment. By 120h, 70%CF treatment had dropped to CK level.



Handle

Total potassium concentration


Total phosphorus concentration

Total nitrogen concentration

Compared with the same time, \*\* was significant

( $p < 0.05$ ), and \*\*\*\* was extremely significant ( $p < 0.001$ ).

Fig. 2 Nutrient content of field drainage at different time after base fertilizer application



Rainfall


Date

Fig. 3 Rainfall of rivers and mountains from June 20 To July 20, 2020

The change of total phosphorus content in ponding was similar to that after base fertilizer application. The peak value of each fertilizer application treatment reached at 1H ( $25.4 \sim 72.2 \text{ mg} \cdot \text{L}^{-1}$ ), which showed that the high fertilizer treatment was significantly higher than the low fertilizer treatment, and it dropped by 70.1%~88.3% at 4h, both significantly higher than CK, and dropped to a very low level at 24h, and then it was  $0.13 \sim 0.57 \text{ mg} \cdot \text{L}^{-1}$  in all treatments.

The change of total potassium content was relatively mild. At 24h, it decreased by 25.0%~52.2% from the peak value, and then changed little at the end of the test, and remained at a high level until the end of the test. In the whole test period, cf and 90% were compared at the same time. There was no same small letter between columns, indicating that the difference between groups was significant ( $p < 0.05$ ). CF treatment was always significantly higher than 80%CF and 70%CF treatment. The above

results showed that reducing the amount of chemical fertilizer could significantly reduce the nutrient content in paddy water, and the decline rate of nutrient content in paddy water was  $p > n > n$ .



Handle

Total potassium concentration

Total phosphorus concentration

Total nitrogen concentration

Fig. 4 Nutrient content of field ponding at different time and different treatment after topdressing

### 2.3 Non point source pollution

Two days after the application of base fertilizer in this test, a heavy rainstorm occurred, and the rainfall in 24h reached more than 150mm (Fig. 3), causing the ponding in the test field to flow out completely. The contents of total nitrogen and total potassium in the ponding in the next day's fertilization treatment decreased rapidly by 94.6% and 94.0%, the same level as CK (Fig. 2). In the early stage of the experiment, the irrigation water was basically kept at about 2cm. This rainstorm caused the loss of total nitrogen and potassium

in the fertilization area to reach 12.0 and 12.7kg·hm<sup>-2</sup> through surface runoff. After removing soil nutrients (CK), the loss of nitrogen and potassium accounted for 16.4% and 15.1% of the base fertilizer application. The phosphorus content in the ponding has dropped to a very low level, indicating that the rainstorm has a great impact on the diversion of nitrogen and potassium nutrients from paddy fields. At the same time, it can be seen from Fig. 2 and Fig. 4 that the nutrient content of paddy water after fertilization decreases with the extension of time, and also decreases significantly with the reduction of fertilization amount, indicating that the amount of fertilization, the time of rainstorm and the interval after fertilization have an impact on the nutrient loss of paddy field. The lower the amount of fertilization and the longer the time after fertilization, the less the nutrient diversion loss of paddy field caused by rainfall, and the lower the risk of pollution to surrounding water bodies.

### 3 Discussion

#### 3.1 Fertilizer reduction and rice production

There is a close relationship between the increase of grain yield and the change of chemical fertilizer application amount. The contribution rate of chemical fertilizer to the yield in China was as high as 56.81%<sup>[2]</sup>. However, due to the habit of relying too much on chemical fertilizer for a long time, the phenomenon of excessive use of chemical fertilizer is widespread, resulting in the rise of production costs, the decline of fertilizer utilization rate, the decline of economic benefits, etc. <sup>[7-9]</sup>. Even in rice, excessive chemical fertilizer (especially nitrogen fertilizer) is easy to cause rice greed at the mature stage, under the condition of high temperature and high heat, a large amount of nutrients accumulated in the plant and could not be transferred to the grain,

but the final yield decreased <sup>[7]</sup>, which verified the truth that "when things reach extremes, they will reverse". In this experiment, the highest rice yield is the conventional fertilization reduction of 10%, and the reduction of 10%~20% on the basis of conventional fertilization has no significant effect on rice yield, which is consistent with zhaoqinglei et al. <sup>[10]</sup> that excessive application of nitrogen and phosphorus fertilizer has no significant effect on rice yield. It shows that the current agricultural main body still has the problem of excessive use of chemical fertilizer, and there is a large space for chemical fertilizer reduction.

#### 3.2 Fertilizer reduction and agricultural non-point source pollution

In order to reduce agricultural non-point source pollution and continuously improve the quality of ecological environment, reducing fertilization is the key work advocated by the Chinese government in agricultural production, and it is also one of the main means to solve the eutrophication of surface water, the high content of nitrate nitrogen in groundwater and the low utilization rate of nutrients. Compared with the conventional fertilization of farmers, appropriately reducing the amount of chemical fertilizer will not only not reduce the yield, but also increase the yield due to improving the population structure of rice, improving the photosynthesis and nutrient transfer capacity of leaves <sup>[11]</sup>, improving the nutrient utilization efficiency of rice and reducing the nutrient loss of rice fields, so as to avoid the risk of water pollution <sup>[5,10,12]</sup>. In this experiment, after applying topdressing for 1h, the contents of total nitrogen, total phosphorus and total potassium in paddy water increased rapidly, and the increase decreased with the decrease of topdressing. Compared with conventional fertilization, the total nitrogen, total phosphorus and total

potassium decreased significantly by 27.1%, 59.5%, 32.7% and 68.2%, 64.8% and 57.5% when the fertilizer was reduced by 20% and 30%. Because the nutrient content in the paddy water was directly related to non-point source pollution [13-14], it showed that reducing fertilization was conducive to reducing the risk of non-point source pollution. Wang et al. [15] investigated the loss of nitrogen in rice field in the application test of different nitrogen amounts. The results showed that the nitrogen loss increased exponentially with the increase of nitrogen application amount. Sun et al. [16] showed through the geographic information system technology scenario analysis that the application amount of nitrogen and phosphorus in rice fields decreased by 10% respectively, and the loss of nitrogen and phosphorus through surface runoff decreased by 7.6% and 3.0% respectively, and believed that the loss of phosphorus was more easily affected by soil properties. After 5 days of balance, the nitrogen content of top dressing in ponding has decreased by 86.1%~91.1%, and the risk of pollution to surrounding water bodies by drainage has decreased significantly; as the soil has a strong phosphorus fixation capacity [13,17], the phosphorus content in the ponding has been in a very low state after 24h (all  $<0.5\text{mg}\cdot\text{L}^{-1}$ ). After 5 days, the 30% reduction treatment is even lower than the class III limit of the surface water quality standard ( $<0.2\text{mg}\cdot\text{L}^{-1}$ ), and there is no risk of non-point source pollution; the potassium content is still maintained at a high level. In case of rainstorm or drainage loss, it is still large. The results showed that the pollution of paddy field ponding nutrient outflow to the surrounding environment was mainly in a period of time after fertilization, and there was a "critical period of pollution risk". Irrigation and drainage before and after the critical period was conducive to greatly reducing the pollution to the surrounding

water environment. Sun et al. [16] also believed that the loss of nutrients in paddy field through surface runoff was mainly within 1 week after fertilization, in which the loss of nitrogen runoff was mainly within 1 week after fertilization, and the loss of phosphorus was mainly within 3~5d, after which the content of nitrogen and phosphorus stabilized at a very low level. Ji et al. [18] found in the investigation of double cropping rice in subtropical hilly and mountainous areas that the high risk period of nitrogen and phosphorus loss is 5 days after the application of base fertilizer and 10~15 days after the application of top dressing. Lentz et al. [19] even believed that the effect of time after fertilization on nutrient loss caused by surface runoff of irrigated fields was greater than the change of fertilization amount. The conclusions of this study are basically consistent with them. However, due to the short time of experimental investigation, it is difficult to confirm whether the contents of nitrogen and potassium other than phosphorus have been reduced to the lowest point, so it is necessary to carry out a further experiment with a longer investigation time.

#### 4 Summary

Under the experimental conditions, the farmers' customary fertilization amount could be reduced by 10%~20% without significantly affecting the rice yield. The suitable fertilization amounts of nitrogen, phosphorus and potassium for rice were 168~189, 72~81 and 120~135 $\text{kg}\cdot\text{hm}^{-2}$  respectively. In the process of rice surface application (topdressing), reducing the amount of conventional fertilizer applied by farmers by 10%~20% can effectively reduce the nutrient content of paddy water, and the total nitrogen, total phosphorus and total potassium decreased by 3.7%~27.1%, 26.3%~59.5% and 5.8%~32.7% respectively. At the same time,

with the extension of time after fertilization, the nutrient content in ponding decreased gradually, and the decreasing rate was phosphorus > nitrogen > potassium. Phosphorus decreased and stabilized at a very low level 24 hours after fertilization, and the risk of environmental pollution was low; the nitrogen content decreased by more than 80% after 5 days, and the pollution risk was significantly reduced. It is suggested that farmers choose the appropriate time for fertilization, try to avoid rainstorm or artificial drainage within 5 days after fertilization, so as to reduce the loss of nutrients in paddy fields and reduce the risk of agricultural non-point source pollution.

## References:

- [1] National Bureau of Statistics. 2019. National data [eb/ol] National Bureau of Statistics, 2019. National data [EB/OL] . [https://data. Stats. Gov. Cn/easyquery. Htm? Cn = C01](https://data.stats.gov.cn/easyquery.htm?cn=C01).
- [2] Ma Kun, diao Gang Study on the contribution rate of chemical fertilizer to the change of grain yield in China [J]. *Journal of plant nutrition and fertilizer*, 2018, 24 (4): 1113-1120
- [3] LIUY Q, WANG H R, JIANG Z M, et al. Genomic basis of geographical adaptation to soil nitrogen in rice [J]. *Nature*, 2021, 590:600-605. [https://doi. Org/101038/s41586-020-03091-w](https://doi.org/101038/s41586-020-03091-w).
- [4] Zhang Zilu, liu Feng, hou Tingyu Current situation of nitrogen and phosphorus loss in paddy fields and research progress of influencing factors in China [J]. *Journal of Applied Ecology*, 2019, 30 (10):
- [5] Lihongjiang, guozhi, jia Jianchu, et al Effects of nitrogen fertilizer reduction on Rice Yield and nitrogen loss in Taihu region [J]. *Journal of ecology*, 2017, 36 (3): 713-718
- [6] Shenjianguo, maweihong, wang Zhong, et al Effects of different slow and controlled release fertilizers on indica Japonica Hybrid Rice [J]. *Zhejiang agricultural science*, 2017, 58 (12): 2153-2156
- [7] QI W, li F R, zhao L, et al. Effects of irrigation and nitrogen application rates on nitrate nitrogen distribution and fertilizer nitrogen loss, wheat yield and nitrogen uptake on a recently reclaimed sandy farmland [J]. *Plant and Soil*, 2010, 337: 325-339.
- [8] SAEED U, wajid S A, khaliq T, et al. Optimizing irrigation and nitrogen for wheat through empirical modeling under semi-arid environment [J]. *Environmental Science and Pollution Research*, 2017, 24:11663 - 11676.
- [9] Shi Changliang, zhujunfeng Economic evaluation and analysis of chemical fertilizer input in grain production in China [J]. *Resources and environment in arid areas*, 2016, 30 (9): 57-63
- [10] Zhaoqinglei, li Yan, ma Jiaqing, et al Effects of different nitrogen and phosphorus levels on Rice Yield and fertilizer absorption and utilization in Nansihu region [J]. *Soil bulletin*, 201142 (5): 1204-1208
- [11] WANG J, fu P H, wang F, et al. Optimizing nitrogen management to balance rice yield and environmental risk in the Yangtze River' s middle reaches [J]. *Environmental Science and Pollution Research*, 2019 (26):4901-4912.
- [12] Yuyingqiu, xuelihong, yanglinzhang Effects of different nitrogen fertilizer management models on nitrogen leakage in paddy soil in Taihu Basin [J]. *Journal of soil*, 2011, 48 (5): 988-995
- [13] QI D L, wu Q X, zhu J Q. Nitrogen and phosphorus losses from paddy fields and the



yield of rice with diferent water and nitrogen management practices [J]. *Scientific Reports*, 2020, 10: 9734.

- [14] LAWLOR P A, helmers M J, baker J L, et al. Nitrogen application rate effect on nitrate-nitrogen concentration and loss insubsurface drainage for a corn-soybean rotation [J]. . *American Society of Agricultural and Biological Engineers*, 2008, 51 (1): 83-94.
- [15] WANG J L, fu Z S, chen G f, et al. Runoff nitrogen (N) losses and related metabolism enzyme activities in paddy field under different nitrogen fertilizer levels [J]. . *Environmental Science and Pollution Research*, 2018, 25: 27583-27593.
- [16] SUN X X, liang X Q, zhang F, et al. A GIS-based upscaling estimation of nutrient runoff losses from rice paddy fields to a regional level [J]. . *Journal of Environmental Quality*, 2016, 45: 1865-1873.
- [17] Xi Bin, zhailimei, liu Shen, et al Effects of combined application of organic and inorganic fertilizers on Maize Yield and soil nitrogen and phosphorus leaching [J]. *Journal of plant nutrition and fertilizer*, 2015, 21 (2): 326-335
- [18] JI L, ouyang X Q, shen J L, et al. Nitrogen and phosphorus runoff losses were influenced by chemical fertilization but not by pesticide application in a double ricecropping system in the subtropical hilly region of China [J]. . *Science of the Total Environment*, 2020, 715: 136852.
- [19] LENTZ R D, lehrsch G A. Nutrients in runoff from a furrow-irrigated field after incorporating inorganic fertilizer or manure [J]. *Journal of Environmental Quality*, 2010, 39: 1402-1415.