

ORIGINAL RESEARCH ARTICLE

Effect of the application of composted manure on the mobility of Cu chemical fractions in a soil contaminated with mining waste

Mayumi Noemí Mendoza-Jiménez¹, Rosa Isela Quintero-Soriano¹, Victor Manuel Duarte Zaragoza^{1*}, Violeta Carrasco-Hernández²

¹ División de Ingeniería Ambiental, Tecnológico de Estudios Superiores del Oriente del Estado de México, Paraje San Isidro, s/n, Mpio. La Paz, Estado de México, 56400, Mexico

² Depto. de Ciencias Forestales, Universidad Autónoma Chapingo, Carretera México-Texcoco, km 38.5, Chapingo, Estado de México, 56230, Mexico. E-mail: *vduarte896@gmail.com

ABSTRACT

The weathering of mining wastes with a high content of metallic sulfides is involved in the release and mobility of heavy metals, being one of the main risk factors for the environment and public health. In this work, two types of manure were used to evaluate their effect on the mobile or bioavailable chemical fractions of Cu in a soil contaminated with mining waste. An experiment was conducted using a soil artificially contaminated with 25% mining waste from Zimapán, to which increasing doses of composted cow and pig manure (0, 3, 6, 12 and 24%) were added. The pseudo-total Cu concentration was determined by atomic absorption spectrophotometry after acid digestion, while the Cu chemical fractions were determined from sequential extractions. The results obtained showed a high pseudo-total Cu concentration in the mining residues and low in the soil and in both types of manure. In the treatments with greater application of pig manure, there was a decrease in the concentration of soluble-interchangeable Cu and an increase in the concentration of Cu strongly bound to the organic fraction. While with cow manure there were higher concentrations of soluble-interchangeable Cu and an increase in the fraction of Cu weakly bound to the organic fraction.

Keywords: soil contamination; environmental analysis; sequential chemical fractionation; contaminated soils; tailings dams

1. Introduction

Mining is one of the most important economic sources in Mexico, generating wealth, but at the

same time causing significant damage to the environment; its operation causes deforestation, habitat destruction, contamination of surface water bodies, and the generation of large quantities of waste^[1]. The latter represent a source of contamination and a public

ARTICLE INFO

Received: May 8, 2022 | Accepted: June 15, 2022 | Available online: July 3, 2022

CITATION

Mendoza-Jiménez MN, Quintero-Soriano RI, Zaragoza VMD, et al. Effect of the application of composted manure on the mobility of Cu chemical fractions in a soil contaminated with mining waste. *Pollution Study* 2022; 3(2): 8 pages.

COPYRIGHT

Copyright © 2022 by author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), permitting distribution and reproduction in any medium, provided the original work is cited.

health problem because they contain metallic sulfides, high concentrations of heavy metals, lack the physical, chemical and biological properties necessary for the development of vegetation, and because they are fine-particle, unstable materials that are dispersed into the environment by water and wind dragging, the latter causing erosion^[2].

The degree of heavy metal contamination in soils is usually measured from the total concentration^[3]. However, this parameter does not provide complete information on ecological risk, geochemical distribution, mobility or bioavailability^[4]. To know the risk posed by heavy metal contamination, it is recommended to quantify the bioavailable concentration, defined as the amount of heavy metal dissolved in soil solution or present in exchange sites^[5]. The soluble and exchangeable forms are considered the most mobile and bioavailable, while those that are incorporated into other components act as a reservoir, remaining retained with organic matter (OM), bound with carbonates and occluded with Fe and Mn oxides or inactive within the crystalline structures of clays^[6].

The mobile concentration of heavy metals is determined from simple extractions with: neutral salts such as CaCl₂ 0.001-0.01 M, NaNO₃ 0.1 M or NH₄NO₃ 1.0 M, and organic chelates such as EDTA and DTPA 0.05 M^[7]; or sequential extractions are performed to quantify the labile fraction of contaminants and their distribution with soil components^[8]. Sequential extractions are procedures applied to evaluate the mobility of heavy metals in sediments, soils and waste materials, and dissolve the different fractions of minerals that retain most of the trace elements, using a series of selective reagents^[9].

Currently, the application of organic amendments is not only an alternative to improve the physical, chemical and nutritional conditions of the soil, but also an option to immobilize soluble heavy metals^[10]. Among the organic materials used for that purpose are composts^[11], biosolids^[12] and animal manures^[13], among others. The amount, composition

and dynamics of soil organic matter (SOM) influence the mobility of heavy metals^[14], as well as other abiotic factors such as pH, Eh, cation exchange capacity, clay content and type, salt content, Fe and Mn oxy-hydroxides, etc.^[15]. Generally, it has been established that humified SOM forms more stable complexes with heavy metals, while fresh OM or light fraction of SOM, can increase the mobility and bioavailability of metals^[16].

The objective of this study was to evaluate the effect of the application of composted cow and pig manure on the mobility of the chemical fractions of Cu in a soil contaminated with mining residues, considering as an analysis variable the proportion of manure applied in the retention and type of manure in the changes of the mobile or bioavailable fractions of Cu. As a working hypothesis, it was proposed that in the treatments with a higher proportion of manure applied, as well as in the manure with a higher cation exchange capacity, a lower concentration of Cu in the fraction with higher mobility (soluble-interchangeable) was recorded.

2. Materials and methods

2.1. Sampling and sample preparation

Sampling of mining waste was carried out on the slopes and upper part of the Santa María deposit in Zimapán, Hidalgo, located at geographical coordinates 0°44'17.6 "N 99°24'03.1" W, and from which a sample representing the Cu concentration of the site was taken, composed in turn of 10 subsamples collected at 10 different points using an auger 30 cm long and 7 cm in diameter, with which 0.5 kg of waste was extracted per subsample. The soil was collected 10 km west of the sampled waste deposit at 20° 42' 10.764 N "99° 20' 5.848" W. The cow and pig manure were composted for four months in the backyard area of the Tecnológico de Estudios Superiores del Oriente del Estado de México and the samples were air dried and dried in the shade at room temperature.

2.2. Experimental design

The experiment was carried out in a greenhouse, based on the preparation of different treatments using cow and pig manure applied in different proportions to the soil artificially contaminated with mining residues (25% w/w). The manure proportions were 0, 3, 6, 12 and 24% (w/w). The experimental design was completely randomized with five treatments and three replicates for each type of manure. The experiment was maintained for two months at an average temperature of 30 °C, humidity was kept at 5% w/w and checked by gravimetry every third day. The total weight of each experimental unit was 1 kg in weight, made from PVC pipes of 35 cm in height and 2.5" in diameter. Changes in the mobile fraction (F1, Soluble-Interchangeable) of Cu were determined from a sequential chemical procedure.

2.3. Analytical methods

The pH was determined by potentiometry^[17] at a soil:water ratio of 1:4, then in the same solution after 20 h of equilibrium the electrical conductivity (EC) was measured with a conductivity bridge (Soil Survey Staff, 1975). The following analyses were determined by different methods: Organic matter (OM) and organic carbon (OC) by the Walkey & Black method modified by Nelson & Sommers^[18]; total nitrogen (TN) by the micro Kjeldahl method and Texture analysis was performed by the Bouyoucos hydrometer technique^[19]; cation exchange capacity (CEC) by the 1N ammonium acetate method pH 7^[20]; pseudo-total Cu content was determined in triplicate after acid digestion with the mixture of HNO₃:HClO₄:H₂O₂ (4:1:1) by Atomic Adsorption Spectrophotometry (AAS) with the Thermo Scientific ICE 3000 Series AA Spectrometer. Calibration was performed using a certified 1,000 mg/L Cu standard (Perkin Elmer Pure: Part N9300183) with a calibration curve in the range of 0.2 - 1.6 mg/L.

2.4. Sequential procedure for the chemical fractions of Cu

The procedure for the determination of the Cu

chemical fractions of each of the treatments consisted of using different chemical reagents sequentially and thus obtaining the fractions in the order described in **Table 1**^[21].

2.5. Statistical analysis

The following analyses were performed: a) analysis of variance (ANOVA) for a distribution of treatments, using the statistical software SAS v9.4 (Statistical Analysis System), and b) Pearson's correlation and Tukey's test for comparison of means.

3. Results and discussion

3.1. Physical and chemical characterization of materials and pseudo-total Cu content.

The mining residues had a sandy texture, with 85% sand, 2.5% clay and 12.5% silt and low MO content, pH of 6.64, EC of 2, 100 dS/m, and low CEC. On the other hand, the soil presented a loam texture, with 14% clay, 48% silt and 38% sand, slightly alkaline pH, EC of 280 dS/m, CEC of 13 Cmol(+)/kg (**Table 2**). Pig manure had a neutral pH; while cow manure recorded a slightly alkaline pH, the latter with higher EC, lower MO content and lower CEC (**Table 2**). On the other hand, the pseudo-total Cu concentration in the mining residues was 2,417 mg/kg, while in the soil it was only 46 mg/kg and 76 mg/kg in the cow manure and 39 mg/kg in the pig manure, the latter registering a higher MO and CO content than the other material.

On the other hand, when both swine and cow manure were applied, changes in the chemical properties of the treatments were observed (**Table 3**): pH increased slightly as the proportion of cow manure increased, but not with swine manure where there was a slight decrease, remaining at a neutral pH. On the other hand, CEC changed from 12.6 to 23 Cmol(+)/kg with pig manure and only up to 14.30 Cmol(+)/kg with cow manure. EC increased from 810 in the 0% manure treatment to 960 dS/m with swine and 1,240 dS/m with cow manure in the 24% treatments.

Table 1. Sequential procedure for the determination of Cu chemical fractions

Step	Fraction	Method and reagents used
F1	Cu-soluble interchangeable	Extraction with acetic acid 1M at pH 5
F2	Cu-bonded to Fe and Mn oxides	Extraction with 0.1 M hydroxylamine hydrochloride and 25% acetic acid.
F3	Cu-bonded weakly with MO	Extraction with HCl 0.1 M.
F4	Strongly Cu-joined with MO	Extraction with 0.5 M NaOH and digestion with 65% HNO ₃ .
F5	Cu-bound to the sulfide phase	Digestion with 8 M HNO ₃ .
F6	Cu-bonded in the residual solid	Digestion with HNO ₃ + HCl, + H ₂ O ₂

Table 2. Physical and chemical characteristics of soil, mining waste and manure

Variable	pH	Texture	CE (dS/m)	CIC (Cmol/kg)	MO (%)	CO (%)	C/N	Total N (%)
Soil	7.89	Franca	280	13	3.3	1.19	-	-
Mining waste	6.64	Arenosa	2100	5.8	1.5	0.87	-	-
Pig manure	6.8	-	789	30	26	17.4	3.6	4.85
Cow dung	7.8	-	1285	12	18	10.44	3.5	3

Table 3. Average chemical parameters measured in the different treatments

Variable	Treatments									
	0%	3%	6%	12%	24%	0%	3%	6%	12%	24%
	Cow dung					Pig manure				
pH	7.1	7.2	7.4	7.7	7.8	7.1	7.1	7.0	7.0	6.9
EC (dS/m)	810	956	1089	1136	1240	810	816	837	924	960
CIC (Cmol(+)/kg)	12.6	12.9	13.4	14.26	14.30	12.6	13.7	18.1	21.3	23.0

Soil components are involved in controlling the mobility of heavy metals, under neutral and alkaline pH conditions there is less mobility and low bio-availability, while acid pH increases their mobility. Soluble salts have an important influence on the mobility of heavy metals in soil^[22] and with the application of manures, these increased, which may influence their mobility. In the case of CEC, it is related to the capacity to exchange metal ions from the soil solution to the surface of clays, Fe and Mn oxides and to the MOS^[23], although the CEC found in the applied manures was lower than in the results of other works with the same materials (up to 36 Cmol(+)/kg have been reported), this variable is related to the type and proportion of OM^[24]. According to Alloway & Trevors^[25] and Caporale & Violante^[26], other soil components that are also responsible for controlling mobility are the concentration and type of clays, SOM, Fe and Mn oxides, carbonates, microorganisms, and organo-mineral complexes. MO participates in controlling the mobility of heavy metals in soil, but it depends on its composition, salt content, C/N ratio, and degree of humification, so it becomes difficult to predict its behavior^[27].

On the other hand, the mining residues presented a high Cu content, with a pseudo-total concentration similar to those reported by Duarte et al.^[28], in the same residues. In addition, it was observed that both the soil and the manure used, being untreated materials, present Cu, although at very low levels of those considered for contaminated soils^[29].

3.2. Chemical fractions of Cu in the treatments

Figure 1 shows the graphs of the chemical fractions of Cu with the application treatments of the two types of manure, where the error bars corresponding to their standard deviation and the Tukey test for the comparison of means are also shown.

In F1 (exchangeable Cu-soluble) a concentration of 22.5, 25.5, 17.8, 28.5 and 18.7 mg/kg Cu was recorded in the respective treatments of 0, 3, 6, 12 and 24% pig manure, with a mean of 22.4 mg/kg Cu. In the cow dung treatments, 33, 28.5, 45.1, 47.6 and 42.4 mg/kg Cu were recorded in the treatments at 0, 3, 6, 12 and 24%, respectively; with a mean of 39.3 mg/kg Cu. In both materials applied with significant difference between treatments ($P < 0.0001$).

In F2 (Cu-bound to Fe and Mn oxides), with pig manure 14.7, 21.7, 24.9, 18.2 and 16.6 mg/kg Cu were obtained in the 0, 3, 6, 12 and 24% treatments, respectively; with a mean of 19.3 mg/kg Cu, with significant difference among treatments ($Pr = 0.0001$). While with cow manure, 4.8, 13.3, 15.7, 17.5 and 10.2 mg/kg Cu were recorded with the application of 0, 3, 6, 12 and 24% manure, respectively; with a mean of 12.3 mg/kg Cu, no significant difference was observed among treatments ($Pr < 0.4672$).

In F3 (Cu-weakly bound to OM), treatments with pig manure application recorded 7.7, 16.4, 11.3, 21.1 and 14.4 mg/kg Cu at 0, 3, 6, 12 and 24% treatments, respectively; with a mean of 2.3 mg/kg Cu and a significant difference among treatments ($Pr < 0.0001$). While in the treatments with cow dung 5.3, 16, 11.2, 21.1 and 14.4 mg/kg Cu in the proportion of 0, 3, 6, 12 and 24%, respectively; with a mean 14.1 mg/kg Cu, and a significant difference between treatments ($Pr < 0.0001$).




Figure 1. Chemical fractions of Cu in pig and cow manure treatments; F1 = Cu-soluble-exchangeable, F2 = Cu-bound to Fe and Mn oxides, F3= Cu-bound weakly to OM, F4 = Cu-bound strongly to OM, F5 = Cu-bound to sulfide phase and F6 = Cu-bound to residual solid. Treatments with the same letter in each graph are not significantly different (Tukey test: $\alpha = 0.05$). Personal elaboration

In F4 (Cu-strongly bound to OM) pig manure treatments, 3.7, 9.4, 10.3, 14.6 and 13.8 mg/kg Cu were found in the 0, 3, 6, 12 and 24% treatments, respectively; with a mean of 14.59 mg/kg Cu. On the other hand, the treatments with cow manure, registered values of 4.1, 2.7, 4.2, 2.4 and 3.1 mg/kg Cu in

the respective treatments of 0, 3, 6, 12 and 24%; with a mean of 3.3 mg/kg Cu. With both materials, a statistical difference between treatments was observed ($Pr < 0.0001$).

In F5 (Cu-bound to the sulfide phase), concentrations of 16.6, 22.6, 24.9, 17.9 and 27.6 mg/kg Cu

were recorded in the respective treatments of 0, 3, 6, 12 and 24%; with a mean of 29.9 mg/kg Cu, with no significant difference among treatments ($Pr = 0.5510$). Very similar concentrations were recorded in the cow manure treatments, presenting 18.2, 25.6, 21.1, 14.6 and 24.9 mg/kg Cu in 0, 3, 6, 12 and 24% manure, respectively; with a mean of 20.9 mg/kg Cu, with no significant difference among treatments ($Pr = 0.2310$).

In F6 (Cu-bound to residual solid), with pig manure application, 32.7, 41.8, 38.9, 48.9 and 49.43 mg/kg Cu were found in the respective treatments of 0, 3, 6, 12 and 24%; a mean of 42.4 mg/kg Cu, with no significant difference between treatments ($Pr = 0.2310$). While in the cow dung treatments, concentrations of 34.8, 58.6, 59.1, 66.8 and 60.1 mg/kg Cu were recorded at 0, 3, 6, 12 and 24% dung, respectively; a mean of 50.9 mg/kg Cu, and significant difference between treatments ($Pr = 0.0105$).

With Pearson's correlation analysis, the following results were obtained from: the concentration of Cu chemical fractions and soil chemical properties (pH, EC and CEC) for each type of manure used. With swine manure, F1 recorded a low correlation coefficient (r), with pH $r = -0.1$, EC $r = -0.1$ and CIC $r = -0.21$; F2 with pH $r = -0.27$, EC $r = -0.32$ and CIC $r = -0.04$; F3 with pH $r = 0.48$, EC $r = 0.29$ and CIC $r = 0.22$; F4 with pH $r = 0.77$, EC $r = 0.84$ and CIC $r = 0.90$; F5 with pH $r = 0.22$, EC $r = 0.37$ and CIC $r = 0.49$; and F6 with pH $r = 0.75$, EC $r = 0.89$ and CIC $r = 0.86$.

With cow manure F1 recorded a high correlation coefficient (r), with pH $r = 0.71$, EC $r = 0.74$ and CIC $r = 0.80$; F2 with pH $r = 0.28$, EC $r = 0.55$ and CIC $r = 0.51$; F3 with pH $r = 0.54$, EC $r = 0.55$ and CIC $r = 0.64$; F4 with pH $r = -0.42$, EC $r = -0.35$ and CIC $r = -0.55$; F5 with pH $r = -0.11$, EC $r = 0.14$ and CIC $r = -0.15$; and F6 with pH $r = 0.59$, EC $r = 0.81$ and CIC $r = 0.76$. In summary, the highest correlations with soil chemical parameters were F4 and F6 in pig manure and F1, F3 and F6 in cow manure.

The different soil mixtures recorded an average

pseudo-total concentration of 664 mg/kg of Cu, a level that, according to USEPA^[30], should be considered as contaminated soil with environmental risk. However, it is relevant to mention that in the experiment the concentration of Cu in F1 (exchangeable soluble) represented the bioavailable fraction for plants and animals, due to its higher mobility. According to Tembo, Sichilongo & Cernak^[31], the World Health Organization (WHO) sets a limit of total Cu concentration in soils of 30 mg/kg to consider it contaminated. In the study conducted by Duarte et al.^[32] they evaluated the effect of cow dung on the immobilization of Pb, Cd, Cu and Zn directly in mining waste, where they reported a distribution of chemical fractions different from those found in this work, possibly because the waste belonged to a different mining waste deposit, although from the same area. In the soluble-interchangeable fraction of Cu, it was expected to register a lower concentration of Cu in the treatments with more manure applied; however, with cow manure F1 increased probably due to an increase in EC by the effect of soluble salts on the mobility of metals in the soil as has been reported by (Usman, Kuzyakov & Stahr^[33] and Acosta, Jansen, Kalbitz, Faz & Martínez-Martínez^[34]). Another possibility could be variation in temperature and humidity, already reported as an influence on the mobility of heavy metals in the soil.

Rodríguez, Ruiz, Alonso-Azcárate & Rincón established an environmental risk code, which is based on the strength of the bond between metals and the different geochemical fractions of the soil, the contaminants contained in the Soluble-Interchangeable Fraction indicate the risk they have on the environment, when F1(soluble-interchangeable), is less than 1%, of the total concentration of the element, there is no risk; therefore, when F1 is in the following ranges the level of risk is valued for example: in a range of 1-10% the risk is low; 11-30% medium; 31-50% high and greater than 50% the risk is very high. What this work indicates is that there is a low risk in the result with the treatments where pig manure was applied, since for F1 it was from 1.3 to 4.7%; and with cow manure it was from 7 to 11%

with a low to medium risk.

SOM has a high influence on the retention of heavy metals, with their composition and dynamics in the soil being important factors in controlling their mobility. However, metals bound to the light fraction of OM (F3) are more susceptible to oxidation and release of contaminants; while metals bound to the heavy fraction of OM, due to the content of humic acids (F4), have a higher stability in the soil. Cow dung presented a higher concentration of Cu in F3, where the light fraction of the OM is extracted, which according to Bloomfield, metal complexes with fulvic acids, can be promoters of mobility in the soil. On the other hand, metals bound to the sulfide phase (S=) after sulfur oxidation (SO =) are released in their cationic phase M⁺⁺ contaminating soils and surface water bodies (Concas, Montinaro, Pisu & Cao). The heavy metals recorded in the residual fraction (F6) are strongly stable, and bound to the crystal lattice of minerals.

4. Conclusions

The high pseudo-total concentration of Cu contained in the mining waste indicates an important risk for the environment and the health of the people and communities surrounding Zimapán; however, with the use of remediation strategies using organic materials, the bioavailable concentrations can be reduced and the specific risk represented by each of the waste deposits in the environment can be determined. A lower concentration of soluble-interchangeable Cu was not observed in the treatments with greater application of cow manure, probably due to the increase of soluble salts, or to the presence of a high proportion of soluble compounds; while with pig manure there was a decrease, but not statistically significant. The application of manure modified the chemical fractions of Cu, in cow manure there was a higher concentration in F3, and in swine manure in F4, Cu strongly bound with OM, it is related to the quality of the organic matter added, which is very important in the processes of Cu retention. The two types of manure seem to have a strong interaction with the soil and mining residues, probably due to

the contribution of clay minerals, since a high correlation is observed between the proportion added and the concentration of copper retained in the residual fractions.

References

1. Higuera P, Oyarzun R. Curso de Minería y Medio Ambiente Medio ambiente Minero. [Internet] Previa.uclm. es. 2002. Available at: <https://previa.uclm.es/users/higuera/mam/>.
2. He Z, Yang X, Soffellab PJ. Trace elements in agroecosystems and impacts on the environment. Review. Journal of Trace Elements in Medicine and Biology 2005; 19: 125-140. doi: 10.1016/j.jtemb.2005.02.010.
3. Rieuwerts JS, Thornton I, Farago ME *et al.* Factors influencing metal bioavailability in soils: Preliminary investigations for the development of a critical loads approach for metals. Chemical Speciation and Bioavailability 1998; 10(2): 61-75.
4. Davutluoglu OI, Seckin G, Ersu CB *et al.* Heavy metal content and distribution in surface sediments of the Seyhan River, Turkey. Journal of Environmental Management 2011; 92: 2250-2259. doi: <http://dx.doi.org/10.1016/j.jenvman.2011.04.013>.
5. Zeien H, Bruemmer GW. Chemical extractions to identify heavy metal binding forms in soils. Mitteilungen der Deutschen Bodenkundlichen Gesellschaft 1989; 59: 505-510.
6. Ahumada I, Mendoza J, Navarrete E *et al.* Sequential extraction of heavy metals in soil irrigated with waste water. Communications in Soil Science and Plant Analysis 1999; 30: 1507-1519. <https://doi.org/10.1080/00103629909370303>.
7. Rog-Young K, Jeong-Ki, Y, Tae-Seung K *et al.* Bioavailability of heavy metals in soils: Definitions and practical implementation-a critical review. Environmental Geochemistry and Health 2015; 37: X-X. doi: 10.1007/s10653-015-9695-y.
8. Vega FA, Covelo EF, Andrade ML. Competitive sorption and desorption of heavy metals in mine soils: influence of mine soil characteristics. Journal of Colloid and Interface Science 2006; 298: 582-592.
9. Gleyzes C, Tellier S, Astruc M. Fractionation studies of trace elements in contaminated soils and sediments: a review of sequential extraction procedures. Trends in Analytical Chemistry 2002; 21: 451-467. DOI: 10.1016/S0165-9936(02)00603-9.
10. Clemente R, Bernal MP. (2006). Fractionation of heavy metals and distribution of organic carbon in two contaminated soils amended with humic acids. Chemosphere 2006; 64: 1264-1273. DOI: 10.1016/j.chemosphere.2005.12.058.
11. Farrell M, Perkins W, Hobbs PJ *et al.* Migration of heavy metals in soil as influenced by compost amendments. Environmental Pollution 2010; 158: 55-64. DOI: 10.1016/j.envpol.2009.08.027.

12. Campos T, Chaer G, Leles P *et al.* Leaching of Heavy Metals in Soils Conditioned with Biosolids from Sewage Sludge. *Floresta e Ambiente* 2019; 26: 2-10. <https://doi.org/10.1590/2179-8087.039918>.
13. Sungur A, Soylak M, Yilmaz S *et al.* Heavy metal mobility and potential availability in animal manure: using a sequential extraction procedure. *Journal of Material Cycles and Waste Management* 2016; 18: 563-572. DOI: 10.1007/s10163-015-0352-4.
14. Zhou T, Wu L, Luo Y *et al.* Effects of organic matter fraction and compositional changes on distribution of cadmium and zinc in long-term polluted paddy soil. *Environmental Pollution* 2018; 232: 514-522. <https://doi.org/10.1016/j.envpol.2017.09.081P>.
15. Rieuwerts JS, Thornton I, Farago ME *et al.* Factors influencing metal bioavailability in soils: Preliminary investigations for the development of a critical loads approach for metals. *Chemical Speciation and Bioavailability* 1998; 10(2): 61-75.
16. Wiatrowska K, Komisarek J. Role of the light fraction of soil organic matter in trace elements binding. *Plos One* 2019; 14(5): e0217077. <https://doi.org/10.1371/journal.pone.0217077>
17. Palmer RG. Laboratory manual. Mexico: Libros y editoriales S.A. D. F. 1979. p. 158.
18. Nelson DW, Sommers LE. Total carbon and organic matter. In: Woodwell, G.M. (ed.) *Methods of soil Analysis Part 3 Chemical Methods*. Soil Science Society of America 1982; 5: 961-1010.
19. Gee GW, Bauder JW. Particle size analysis. In Klute, A. (ed.) *Methods of soil Analysis. Part 1. Physical and Mineralogical Methods*. 2nd Edition. Agronomy Monograph 9. ASA and SSSA, Madison, WI. 1986. p. 404- 407.
20. Rowell DL. *Soil Science: Methods and Applications*. 1994. Longman, Harlow.
21. Pagnanelli F, Mosca E, Giuliano V *et al.* Sequential extraction of heavy metals in river sediments of an abandoned pyrite mining area; pollution detection and affinity series. *Environmental Pollution* 2004; 132: 189-201. DOI: 10.1016/j.envpol.2004.05.002.
22. Hund-Rinke K, Koerdel W. Underlying issues in bioaccessibility and bioavailability: Experimental methods. *Ecotoxicology and Environmental Safety* 2003; 56: 52-62.
23. Sparks DL. *Environmental soil chemistry*. 2nd ed. 2002. San Diego: Academic.
24. Alloway BJ, Trevors JT. *Heavy metals in soils: Trace metals and metalloids in soils and their bioavailability*: Dordrecht, Springer. 2013. 613 p.
25. Caporale A, Violante A. Chemical Processes Affecting the Mobility of Heavy Metals and Metalloids in Soil Environments. *Current Pollution Reports* 2016; 2: 15-27. doi: 10.1007/s40726-015-0024-y.
26. Shi W, Lu C, He J *et al.* Nature differences of humic acids fractions induced by extracted sequence as explanatory factors for binding characteristics of heavy metals. *Ecotoxicology Environmental Safety* 2018; 154(15): 59-68. <https://doi.org/10.1016/j.ecoenv.2018.02.013> PMID: 29454987.
27. USEPA. *Soil Screening Guidance. Technical Background*. Document. U.S. Gov. Print. Office. 1996. Washington, D. C. USEPA Rep.504/R-95/128.
28. Tembo BD, Sichilongo K, Cernak J. Distribution of copper, lead, cadmium and zinc concentrations in soils around Kabwe town in Zambia. *Chemosphere* 2006; 63: 497-501. doi: 10.1016/j.chemosphere.2005.08.002.
29. Duarte VM, Carrillo-González R, Lozano ML *et al.* Fractionation of heavy metals in mine tailings amended with composted manure. *Soil and Sediment Contamination: An International Journal* 2019; 28(2): 148-161. doi: 10.1080/15320383.2018.1553931.
30. Usman AR, Kuzyakov Y, Stahr K. Effect of immobilizing substances and salinity on heavy metals availability to wheat grown on sewage sludgecontaminated soil. *Soil & Sediment Contamination* 2005; 14: 329-344.
31. Abollino O, Aceto M, Malandrino M *et al.* Distribution and mobility of metals in contaminated sites. *Chemometric Investigation of Pollutant Profiles. Environmental Pollution* 2002;119: 177-193. doi: 10.1016/s0269-7491(01)00333-57.
32. Hayes (Editors). Bloomfield C. *The translocation of metals in soils*. In: D.J. Greenland and M.H.B. *The Chemistry of Soil Processes*. John Wiley & Sons Ltd. 1981. Chichester.
33. Concas A, Montinaro S, Pisu M *et al.* Mechanochemical remediation of heavy metals contaminated soils: Modelling and experiments. *Chemical Engineering Science* 2007; 62(18.20): 5186-5192. <https://doi.org/10.1016/j.ces.2007.02.024>.
34. Jiang M, Zeng G, Zhang C. Assessment of heavy metal contamination in the surrounding soils and surface sediments in Xiawangang River, Qingshuitang District. *Plos One* 2013; 8(8): e71176. doi: 10.1371/journal.pone.0071176.