

Article

Recovering of soil contaminated by hydrocarbons mixing

Juna Manuel Sánchez-Yáñez

Environmental Microbiology Laboratory, Research Institute (Chemical Biological, Chemical and Biological), Michoacán University of San Nicolás de Hidalgo, Morelia 58000, Mexico; syanez@umich.mx

CITATION

Sánchez-Yáñez JM. Recovering of soil contaminated by hydrocarbons mixing. *Pollution Study*. 2023; 4(2): 2014.
<https://doi.org/10.54517/ps.v4i2.2014>

ARTICLE INFO

Received: 19 July 2023
Accepted: 10 August 2023
Available online: 20 August 2023

COPYRIGHT

Copyright © 2023 by author(s).
Pollution Study is published by Asia Pacific Academy of Science Pte. Ltd. This work is licensed under the Creative Commons Attribution (CC BY) license.
<https://creativecommons.org/licenses/by/4.0/>

Abstract: In Mexico, an agricultural soil poor in nitrogen (N) contaminated by a hydrocarbon derivative such as automotive residual oil (ARA), with a relatively high concentration of 100,000 ppm, is an environmental problem, but also because it drastically affects soil properties associated with the mineralization of organic matter and loss of fertility, since it exceeds the maximum accepted limit of 4400 ppm of the Mexican standard called, NOM-138-SEMARNAT-2012 (NOM-138). An alternative solution is to treat it with ecological actions to eliminate the ARA and recover fertility. Therefore, the objectives of this research were: i) bioremediation of soil contaminated by 100,000 ppm of ARA ii) phytoremediation using *Sorghum vulgare* with *Aspergillus inger* and *Penicillium chrysogenum* to decrease ARA to a value below 4400 ppm of NOM-138. For this purpose, soil recovery was performed using the variable-response: disappearance of ARA by Soxhlet at the beginning and after bioremediation and at the end of phytoremediation with *S. vulgare* with phenology and biomass to seedling. All experimental data were validated by ANOVA/Tukey HSD $P < 0.05\%$. The results indicated that bioremediation and phytoremediation of soil contaminated by 100,000 ppm of ARA, decreased it to 3400 ppm, a value lower than the maximum established by NOM-138, sufficient for soil recovery in agricultural production, in 120 days, a relatively short period of time.

Keywords: soil; ARA; biostimulation; NOM-138; *S. vulgare*; mushrooms

1. Introduction

Currently, some petroleum derivatives, such as oils used in the lubrication and refrigeration of automobiles, generate products that pollute the environment, such as automotive waste oil (ARA), a combination of aliphatic, aromatic and polycyclic hydrocarbons [1]. In Mexico ARA, according to the General Law of Ecological Balance and Environmental Protection [2] is a toxic environmental waste. In order to determine the damage caused by ARA in the soil, there is a Mexican standard [3] known as NOM-138-SEMARNAT-2012 (NOM-138), which establishes the maximum permissible concentration limit, especially for an agricultural soil of 4400 ppm, an amount that prevents the mineralization of organic matter (OM), prevents gas exchange, which consequently decreases or cancels agricultural production [4], given the phytotoxicity of ARA aromatics. The literature reports that a soil impacted by ARA is reduced by chemical methods, which are fast, of high economic value, but which cause collateral damage by leaving toxic residues for animal and/or plant life [5,6]. An alternative ecological solution is bioremediation (BIO) which, by enriching the soil with basic minerals N (nitrogen), P (phosphorus), K (potassium), rebalances the carbon:nitrogen ratio (C:N) caused by excess carbon: nitrogen (C:N):N caused by the excess carbon in the ARA, to eliminate it in a relatively long or short time [7], this speed is dependent on the complexity and concentration of the

hydrocarbons it contains, especially when the soil is poor in N, so it is necessary to enrich it with an animal fertilizer, such as vermicompost (LC), which in addition to the high content of essential minerals, incorporates microorganisms that favor the oxidation of ARA, for this type of BIO, it is important to have an adequate demand of O₂ (oxygen) to ensure the constant elimination of ARA, without causing drastic changes in pH, since neutrality accelerates the oxidation of ARA [8].

In a soil contaminated by 100,000 ppm of ARA, BIO is insufficient to reduce it to a value lower than 4,400 ppm, which allows agricultural recovery in accordance with NOM-138 [3], consequently, phytoremediation (FITO) is indispensable, with plants whose root system tolerates phyto-toxicity to hydrocarbons and facilitates the oxidation of ARA [9–12]. As it is reported for other petroleum products, mainly because these plants can improve the elimination capacity of these hydrocarbons, mainly when inoculated with microorganisms that naturally hydrolyze aromatics. In soil impacted by a relatively high concentration of hydrocarbons: so that, by oxidizing them, it is possible to recover fertility for agricultural production, according to some environmental regulation in force [3]. In relation to improving the capacity of a plant to mineralize soil hydrocarbons, it is reported that genera and species of fungi such as *Aspergillus niger* and *Penicillium chrysogenum* not only stimulate the growth of the plant root system [1,9,13], but also metabolize hydrocarbons similar to those detected in the ARA and consume them until they are reduced to a value that facilitates the recovery of useful soil for agricultural production [14–17]. Based on this information, the objectives of this research were: i) bioremediation of soil contaminated by 100,000 ppm of ARA, ii) phytoremediation with *Sorghum vulgare* inoculated with *A. niger* and *P. chrysogenum* to reduce ARA to a value below the maximum of NOM-138 as evidence of its recovery.

2. Materials and methods

This research was conducted in the greenhouse of the Environmental Microbiology laboratory of the Instituto de Investigaciones Químico Biológicas (IIQB) of the Universidad Michoacana de San Nicolás de Hidalgo (UMSNH). Under the following microclimatic conditions: temperature of 23.2 °C, luminosity of 450 $\mu\text{mol m}^{-2} \text{s}^{-1}$, relative humidity of 67%. It was used, an agricultural soil collected in an area at 19° 37' 10" north latitude and 101° 16' 41.00" west longitude, with an altitude of 2013 masl, with a temperate climate of a place called "Uruapilla" of the municipality of Morelia, Michoacán, Mexico, on the Morelia-Pátzcuaro highway, Michoacán. The ARA was collected from an oil change shop in Morelia, Michoacán, Mexico.

Table 1. Physicochemical properties of agricultural soil not artificially contaminated by automotive waste oil.

Parameter	Interpretation value
pH (1:20)	6.02 acid
Total nitrogen (%)	0.11 poor
Organic matter (%)	0.58 poor
Ion exchange capacity (Cmol ⁽⁺⁾ kg ⁻¹)	26.64 saline
Texture (%) ⁺	50 (Ac) low aeration, 7(L), 43 (Ar)
True density or RD (g/cm ³)	2.22
Bulk density or DA (g/cm ³)	1.05
Porosity (%)	47.5
Field capacity (%)	54 low humidity

⁺Ar: sand, L: silt, AC: clay, *for soils of volcanic origin, **calculated from DA and DR.

**estimated texture, +for clayey soil according to Mexican standard: NOM-021-RECNAT-2000.

Table 1, details the physicochemical properties of the agricultural soil uncontaminated by ARA [18], with a pH of 6.02 slightly acidic, with low MO content 0.58%, a poor concentration of total N of 0.11%, a high ion exchange capacity of 26.64 Cmol⁽⁺⁾ kg⁻¹ or saline, with a texture composition: clay 50 %, silt 7% and sand 43%, so the soil was classified as clayey, this soil was sieved with a No. 20 mesh, solarized for 48 h and reduced the problem of pests and diseases, then contaminated with ARA, and began the BIO of the soil by dissolving the 100,000 ppm of ARA in the commercial detergent “La Corona” at 0.5 % (w/v). Then 1.0 kg of this soil contaminated with ARA was placed in the upper part of the Leonard jar, in the lower part water or mineral solution (SM) was deposited, both parts were connected by a cotton strip, for the movement of the liquid by capillarity (**Figure 1**).




Figure 1. Leonard pitcher diagram [13].

The experiment was divided into i) soil BIO by 100,000 ppm ARA, according to **Table 2**, ii) soil PHYTO by *S. vulgare* with *A. niger* and/or *P. chrysogenum*, according to **Table 3**. With a randomized block experimental design of six

treatments and six replicates: soil without ARA irrigated only with water or absolute control (AC), soil fed with a mineral solution referred to as relative control (RC), soil with ARA without biostimulation or FITO or negative control (NC), and soil with ARA, biostimulated and phytoremediated.

Table 2. Experimental design of bioremediation of agricultural soil contaminated by 100,000 ppm of residual automotive oil.

Agricultural land	100,000 ppm of ARA	Detergent at 0.5% and LC at 3%.	SM at 100% and H ₂ O ₂ at 0.05%.
Absolute control (AC)	-	-	-
Relative control (RC)	-	-	+
Negative control (NC)	+	-	-
Bioremediation	+	+	+

*n = 6; aggregate (+); not aggregate (-). LC vermicompost, SM mineral solution.

In the first phase, the soil with 100000 ppm of ARA was dissolved in the commercial detergent “La Corona” at 0.5% and a LC at 3% for 30 days, then biostimulated with a SM with the following composition (g·L⁻¹): NH₄ NO₃ 10.0, K₂HPO₄ 2.5, KH₂ PO₄ 2.0, MgSO₄ 0.5, NaCl 0.1, CaCl₂ 0.1, FeSO₄ 0.01 and 1.0 mL/L of a microelement solution (g·L⁻¹): H₃BO₃ 2.86, ZnSO₄ 7H₂ O 0.22, MgCl₂·7H₂ O 1.8, adjusted to pH 6.8 [19], was biostimulated with the SM, and simultaneously with H₂ O₂ at 0.05% for 72 h for one month, while to facilitate gas exchange the moisture was adjusted to 80 % of the field capacity of the agricultural soil. The variable-response of soil recovery was ARA detected by Soxhlet at the beginning and end of BIO and FITO [15].

Table 3. Experimental design for phytoremediation of a soil impacted by automotive waste oil remaining from biostimulation.

Sorghum vulgare*.	ARA	A. niger	P. chrysogenum
Irrigated with water (absolute control)	-	-	-
Fed with mineral solution (relative control)	-	-	+
	+	+	-
Biostimulation	+	-	+
	+	+	+

*n = 6; added (+); not added (-), ARA automotive waste oil.

In the second phase, soil contaminated by 100,000 ppm ARA after biostimulation was phytoremediated according to **Table 3** by planting *S. vulgare* obtained from the Secretaria de Agricultura Ganadería y Desarrollo Rural Pesca y Alimentación of the Mexican government. *S. vulgare* was treated with *A. niger* and/or *P. chrysogenum*, both fungi were isolated from decaying wood for their ability to degrade lignin, and molecularly identified as the species indicated 20, to inoculate on *S. vulgare* seeds. They were replicated on avocado pit agar with the following composition (g L⁻¹): avocado pit 10, casein peptone 5, yeast extract 1.3, K₂ HPO₄ 0.17, KH₂ PO₄ 2.61, MgSO₄ 1.5, NaCl 0.9, CuSO₄ 0.05, bromothymol blue 10 ppm, 2.5 mL 10% deter- people solution 1.0 mL trace element solution, agar 18.0 g, pH adjusted to 5.5. *S. vulgare* seeds were disinfected with 0.2 % NaOCl/5 min,

washed 6 times with sterile distilled water, then for every 10 *S. vulgare* seeds were inoculated with 1.0 mL of *A. niger* and/or *P. chrysogenum* equivalent to 1×10^6 Pro-pule-forming units (PFU)/1 mL [19] to be sown in agricultural soil artificially contaminated by ARA, 60 days later, phenology: plant height (PA) and root length (RL), and biomass: aerial and root fresh weight (AFW/RFW) and aerial and root dry weight (AFW/RDW) were measured [19]. Experimental data were validated with ANOVA/Tukey HSD $P < 0.05$ % with the statistical program Statgraphics Centurion [20,21].

3. Results

In **Table 4**, BIO reduced ARA from 100,000 to 37620 ppm ARA in 60 days, a statistically different numerical value relative to 90,000 ppm ARA in soil without bioremediation or CN.

Table 5 shows the phenology of *S. vulgare* boosted with *A. niger* and *P. chrysogenum* at 60 days in soil with 37620 ppm of ARA, 29.2 cm PA, 17.0 cm LR were recorded, both numerical values with statistical difference compared to 20.0 cm PA and 8.5 cm LR of *S. vulgare* without inoculation irrigated only with water in soil without ARA referred to as CA. While *S. vulgare* with *A. niger* and *P. chrysogenum* in soil with ARA, recorded 10.0 g PFA and 5.0 g PFR, both statistically different numerical values compared to 4.77 g PFA and 2.89 g PFR of *S. vulgare* fed with a SM, in soil without ARA used as CR. Regarding the biomass of *S. vulgare* with *A. niger* and *P. chrysogenum* in soil with ARA, 2.2 g of PSA and 1.6 g of PSR were recorded, these numerical values were statistically different compared to the 1.14 g of PSA and 1.31 g of PSR of *S. vulgare* fed with the SM or CR.

Table 4. In soil concentration of residual automotive oil remaining from bioremediation for 60 days.

*Agricultural soil artificially polluted by 100,000 ppm of ARA	Final concentration
Irrigated with water or CN	90000b***
**Multiple Bioremediation	37620a

*B ioremediation: detergent at 0.5%, vermicompost at 3%. Mineral solution at 100% and H₂ O₂ at 0.05%. ***Different letters indicate that they are statistically different according to ANOVA/Tukey at 0.05%.

Table 5. Phenology and biomass of *Sorghum vulgare* with *A. niger* and *P. chrysogenum* after soil phytoremediation with 37,620 ppm of residual automotive oil, after 60 days.

Sorghum vulgare*.	AP (cm)	LR (cm)	PF (g)		ps (g)	
			Aerial	Radical	Aerial	Radical
Irrigated with water (absolute control)	20.0 ^b **	8.5 ^d	5.20 ^c	2.90 ^c	1.15 ^c	1.20 ^c
Fed with mining solution to the	25.0 ^a	16.0 ^b	4.77 ^c	2.89 ^c	1.14 ^c	1.31 ^b
100% (relative control) with <i>Aspergillus niger</i>	18.0 ^c	14.0 ^{bc}	7.27 ^b	3.71 ^b	1.84 ^a	1.28 ^b
with <i>Penicillium chrysogenum</i>	22.0 ^b	15.0 ^b	8.93 ^{ab}	5.73 ^a	1.50 ^b	0.99 ^d
with <i>A. niger</i> and <i>P. chrysogenum</i>	29.2 ^a	17.0 ^a	10.0 ^a	5.0 ^a	2.2 ^a	1.60 ^a

*n = 6. **Different letters indicate statistical difference according to ANOVA/Tukey at 0.05%. AP plant height, LR root length, PF fresh weight, PS dry weight, PS dry weight.

Table 6 shows the decrease in soil ARA from 37,620 ppm to 3400 ppm by the activity of *S. vulgare* with *A. niger* and *P. chrysogenum*, this last value was statistically different compared to the 80000 ppm ARA of the soil without bioremediation and phytoremediation used as CN.

Table 6. In soil concentration of residual automotive oil after phytoremediation with *Sorghum vulgare* enhanced with *Aspergillus niger* and *Penicillium chrysogenum* at 120 days.

*Agricultural soil contaminated by ARA	ARA+ (ppm)	
	Initial	Final
Soil without phytoremediation or negative control	100000a**	80000b
Phytoremediated soil, <i>S. vulgare</i> + <i>A. niger</i> + <i>P. chrysogenum</i>	37620a	3400b

*n = 6; **Different letters are statistically different at 0.05% according to Tukey, + ARA automotive residual oil.

4. Discussion

In **Table 4**, the agricultural soil with 100,000 ppm of ARA, started the BIO with the detergent, which solubilized it, to facilitate the native heterotrophic aerobic microorganisms to carry out a partial reduction of the concentration [4,5,7,8], while the BIO with the LC by enriching the soil with urea allowed the equilibrium of the ratio C:N ratio, so that the native microorganisms could partially oxidize the ARA [22,23], in the same way that the SM with salts NH_4^+ , NO_3^- and PO_4^{-3} accelerated the mineralization of the ARA [24,25], simultaneously the BIO with H_2O_2 supplied the demand O_2 to optimize the oxidation of ARA [14,26], therefore the field capacity of the soil was controlled at 80%, which allowed the exchange of gases and the decrease of the concentration of ARA [17,27] to a sufficient level for the sowing of *S. vulgare* inoculated with *A. niger* and *P. chrysogenum*, in the soil recovery route according to NOM-138.

In comparison with the soil with 100,000 ppm of ARA, used as CN, in which natural attenuation was insufficient to eliminate it, due to the excess of C in the ARA that formed a hydrophobic film of the ARA and prevented the exchange of gases such as O_2 while the lack of minerals essential for the oxidation of ARA prevented that concentration from being reduced [25,26,28].

Table 5 shows the phenology and biomass of *S. vulgare* with *A. niger* and *P. chrysogenum*, planted in soil when the ARA concentration was decreased to 37620 ppm. Where indirectly a decrease in ARA was recorded due to the healthy growth of *S. vulgare* partly because it is naturally tolerant to phytotoxic stress of ARA, and due to the positive effect of *A. niger* and *P. chrysogenum* in the rhizosphere of the plant, where these fungi can generate plant growth-promoting substances [25,29], to enhance the root mineral uptake capacity and decrease the concentration of ARA leading to the recovery of soil health and allowing it to be used in agricultural production [10], in stark contrast to *S. vulgare* uninoculated with *A. niger* and *P. chrysogenum* planted in soil contaminated by 86,000 ppm ARA without bioremediation, where the relatively high concentration caused inhibition of healthy *S. vulgare* growth [11].

Table 6 shows the concentration of soil ARA impacted by 37,620 ppm of ARA remaining from BIO, then by FITO with *S. vulgare* enhanced with *A. niger* and *P. chrysogenum* that stimulated healthy root growth. *chrysogenum* that stimulated healthy root growth, suggests the conversion of root exudates into phytohormones [11,17,30] that increased the amount of root hairs, to expand the area of exploration and mineral uptake, and thereby increased the tolerance of *S. vulgaris* to ARA [10]. In addition, there is evidence that both *A. niger* and *P. chrysogenum* have the capacity to degrade aromatics of ARA [12,14,30,31], and facilitated the elimination of ARA up to a concentration of 3400 ppm, a value lower than the maximum established by NOM-138, thus achieving soil recovery for reuse in agricultural production.

Ethical approval: The approval of the research by the Ethics Committee of the Universidad Michoacana de San Nicolás de Hidalgo—Mexico, followed the guidelines established for this committee.

Conflict of interest: The author declares no conflict of interest.

References

1. Wu M, Dick WA, Li W, et al. Bioaugmentation and biostimulation of hydrocarbon degradation and the microbial community in a petroleum-contaminated soil. *International Biodeterioration & Biodegradation*. 2016; 107: 158-164. doi: 10.1016/j.ibiod.2015.11.019
2. General law of ecological balance and environmental protection. Available online: http://www.diputados.gob.mx/LeyesBiblio/pdf/1_48_240117.pdf (accessed on 2 June 2023).
3. Norma Oficial Mexicana NOM-138-SEMAR NAT/SSA1-2012, Maximum permissible limits of hydrocarbons in soils and guidelines for sampling in the characterization and specifications for remediation. DOF Secretary of the Interior (Spanish). Available online: http://www.dof.gob.mx/nota_detalle.php?codigo=5313544&fecha=10/09/2013 (accessed on 2 June 2023).
4. Thapa B, KC AK, Ghimire A. A Review on bioremediation Of Petroleum Hydrocarbon Contaminants in Soil. *Kathmandu University Journal of Science, Engineering and Technology*. 1970; 8(1): 164-170. doi: 10.3126/kuset.v8i1.6056
5. Shahi A, Aydin S, Ince B, et al. Evaluation of microbial population and functional genes during the bioremediation of petroleum-contaminated soil as an effective monitoring approach. *Ecotoxicology and Environmental Safety*. 2016; 125: 153-160. doi: 10.1016/j.ecoenv.2015.11.029
6. Rivera Ortiz P, Rivera Lárraga JE, et al. Biostimulation and bioremediation of oil-contaminated drill cuttings (Spanish). *Revista Internacional de Contaminación Ambiental*. 2018; 34(2): 249-262. doi: 10.20937/rica.2018.34.02.06
7. Torri SI, Cabrera MN, Alberti C. Potential respiration during biostimulation of a soil contaminated with polycyclic aromatic hydrocarbons. *Rev Int Contam Ambient*. 2018; 34(1):127-36. doi: 10.20937/rica.2018.34.01.11.
8. Reyes Reyes MA, Puentes Cala EA, Casanova Montes EL, López Deluque F, Panqueva Alvarez JH, Castillo Villamizar GA. Immobilization of potentially crude oil-degrading bacteria in natural and synthetic organic matrices. *Rev Int Contam Ambie*. 2018; 34(4): 597- 609. doi: 10.20937/RICA.2018.34.04.04
9. Guevara Espinosa MaD, Cruz Miranda N, Rivera Morales C, et al. Phytoremediation of soils contaminated with Mn and Cu from *Octmum basilicum*. *Rev Latinoam Ambient Cienc*. 2018; 9(22): 6-89.
10. Leitão AL, Enguita FJ. Gibberellins in *Penicillium* strains: challenges for endophyte-plant host interactions under salinity stress. *Microbiol Res*. 2016; 183: 8-18. doi: 10.1016/j.micres.2015.11.004
11. Solyman SNED, Abdel-Monem MO, Abou-Taleb KA, et al. Production of Plant Growth Regulators by Some Fungi Isolated under Salt Stress. *South Asian Journal of Research in Microbiology*. Published online March 4, 2019: 1-10. doi: 10.9734/sajrm/2019/v3i130076
12. Bilal L, Asaf S, Hamayun M, et al. Plant growth promoting endophytic fungi *Asprgillus fumigatus* TS1 and *Fusarium proliferatum* BRL1 produce gibberellins and regulates plant endogenous hormones. *Symbiosis* 2018; 76(2):117-27. doi:

- 10.1007/ s13199-018-0545-4
13. Garcia González MM, Farías Rodríguez R, Peña Cabriales JJ, et al. Inoculation of wheat var. Pavón with *Azospirillum* spp. and *Azoto- bacter beijerinckii*. *Terra Latinoam*. 2005; 23(1): 65-72.
 14. Contreras H, Carreño C. Efficiency of petroleum hydrocarbon biodegradation by filamentous fungi isolated from contaminated soil. *Rev de Investig Agroproduccion Sustentable*. 2018; 1(1): 27-33. doi: 10.25127/ ucn.v1i1.269
 15. Hernández Valencia I, Mager D. Use of *Panicum maximum* and *Brachiaria brizantha* to phytoremediate soils contaminated with a light petroleum crude oil. *Bioagro*. 2003; 15(3): 149-56.
 16. Delgadillo López AE, González Ramírez CA, Prieto García F, et al. Phytoremediation: an alternative to eliminate contamination. *Trop Subtrop Agroecosyst*. 2011; 14(2): 597-612.
 17. Ite AE, Ibok UJ. Role of plants and microbes in bioremediation of petroleum hydrocarbons contaminated soils. *Int J Environ Bioremediat Biodegrad*. 2019; 7(1):1-19. doi: 10. 12691/ijebb-7-1-1
 18. Mexican Official Norm N OM-021-SEMARNAT-2000, which establishes the specifications of fertility, salinity and soil classification, study, sampling and analysis. Available online: http://diariooficial.gob.mx/nota_detalle.php?codigo=717582&fecha=31/12/2002 (accessed on 2 June 2023).
 19. Sánchez-Yáñez J. Brief Treatise on Agricultural Microbiology, theory and practice, Universidad Michoacana de San Nicolás de Hidalgo, Centro de Investigaciones y Desarrollo del Estado de Michoacán (Spanish). Sedagro Cosustenta, SA de CV, Morelia, Michoacan; 2007. p. 118-119.
 20. Baltierra Trejo E, Marquez Benavides L, Sanchez-Yáñez JM. Inconsistencies and ambiguities in calculating enzyme activity: The case of *laccase*. *J Microbiol Methods*. 2015; 119: 126-31. doi: 10.1016/j.mimet.2015.10.007
 21. Walpole ER, Myers RH, Myers SL. Probability and Statistics for Engineering and Science. Available online: <http://librosenpdf.org/libro-pdf-probabilidad-y-estadistica/> (accessed on 2 June 2023).
 22. Riojas González HH, Gortáres Moroyoqui P, Mondaca Fernández I, et al. Influence of surfactants in the remediation of hydrocarbon-contaminated soils. *Bistua*. 2016; 7(1): 94-115. doi: 10.18359/ rfb.2066
 23. Ramos Oseguera CA, Castro Ramírez AE, León Martínez NS, et al. Vermicomposting to recover sandy loam soil fertility and peanut (*Arachis hypogaea* L.) yield. *Terra Latinoam*. 2019; 37(1): 45-55. doi: 10.28940/tl.v37i1.331
 24. Jiménez Hernández V, Guerra Sánchez R. Obtaining an enriched medium to make the bioavailability of weathered hydrocarbons in a coastal soil more efficient. *Rev Int Contam Ambient*. 2016; 32(4): 413-24. doi: 10.20937/RICA.2016.32.04.05.
 25. Alvaro CES, Arocena LA, Martínez MA, et al. Aerobic biodegradation of hydrocarbon fractions from oil activity in a soil of the Northern Patagonia region, Argentina. *Rev Int Contam Ambie*. 2017; 33(2): 247-57. doi: 10. 20937/RICA.2017.33.33.02.06
 26. Velásquez Arias JA. Soil and water contamination by hydrocarbons in Colombia (Spanish). *Análisis de la fitorremediación como estrategia biotecnológica de recuperación*. *Revista de Investigación Agraria y Ambiental*. 2017; 8(1): 151-167. doi: 10.22490/21456453.1846
 27. Mohsenzadeh F, Chehregani Rad A, Akbari M. Evaluation of oil removal efficiency and enzymatic activity in some fungal strains for bioremediation of petroleum-polluted soils. *Iranian Journal of Environmental Health Science & Engineering*. 2012; 9(1). doi: 10.1186/1735-2746-9-26
 28. Barrios Ziolo LF, Robayo Gómez J, Prieto Cada-vid S, et al. Bioremediation of soils contaminated with used motor oils. *Revista Cintex*. 2015; 20(1): 69-96.
 29. Effendi AJ, Kamath R, McMillen S, et al. Strategies for Enhancing Bioremediation for Hydrocarbon-Impacted Soils. In: *Proceeding of the Society of Petroleum Engineers International. Asia Pacific Health, Safety, Security, Environment and Social Responsibility Conference*; 2017. doi: 10.1109/ITME.2015.163
 30. Leitão AL. Potential of *Penicillium* Species in the Bioremediation Field. *International Journal of Environmental Research and Public Health*. 2009; 6(4): 1393-1417. doi: 10.3390/ijerph6041393
 31. Chaudhary S, Shankar A, Singh A, et al. Usefulness of *Penicillium* in enhancing plants resistance to abiotic stresses: An overview. In: Chaudhary S, Shankar A, Singh A, et al. (editors). *New and Future Developments in Microbial Biotechnology and Bioengineering*. Elsevier; 2018. pp. 277-84. doi: 10.1016/ B978-0-444-63501-3.00017-X